

# Surgical and functional outcome of lateral canthal tarsorrhaphy in unilateral paralytic lagophthalmos

Kavita A Dhabarde<sup>1\*</sup>, A H Madan<sup>2</sup>

<sup>1</sup>Assistant Professor, <sup>2</sup>Professor and HOD, Department of Ophthalmology, Government Medical College Nagpur, Maharashtra, INDIA.  
Email: [drkavitad1423@yahoo.com](mailto:drkavitad1423@yahoo.com)

## Abstract

Ten patients having unilateral infranuclear 7th nerve palsy with lagophthalmos of different age group and different aetiology underwent lateral canthal tarsorrhaphy to prevent progression of and treatment of severe exposure keratitis in tertiary eye care centre at Govt. Medical College and Hospital, Nagpur. Patients were advised local lubricants, antibiotics. Nearly all pts showed improvement in lid closure, epiphora and keratitis. Lat. tarsorrhaphy supported the lower lid with not much cosmetic disfigurement of the eyeball. Only in a single case the ulcer perforated. Lateral tarsorrhaphy is a good option for poor patients having infranuclear Bell's palsy with severe exposure keratitis who cannot afford to go for temporalis transplant.

**Keywords:** lateral canthal tarsorrhaphy, unilateral paralytic lagophthalmos.

## \*Address for Correspondence:

Dr. Kavita A. Dhabarde, Flat no. 403, Ganpati apartment, Wanjarinagar, Plot no. 28, Near Samrudhi Co Op Bank, Nagpur, Maharashtra, INDIA.

Email: [drkavitad1423@yahoo.com](mailto:drkavitad1423@yahoo.com)

Received Date: 07/10/2014 Accepted Date: 16/10/2014

## Access this article online

Quick Response Code:



Website:

[www.statperson.com](http://www.statperson.com)

DOI: 16 October 2014

## INTRODUCTION

Facial nerve palsy is common problem that involves the paralysis of any structure innervated by facial nerve. The most common cause of facial nerve palsy is Bell's palsy, an idiopathic disease that may only be diagnosed by exclusion. Exposure keratitis occurs in facial nerve palsy and may lead to visual loss from corneal damage unless it is treated appropriately. To prevent and / or treat exposure keratitis and reestablish eyelid function tarsorrhaphy is done

## AIMS AND OBJECTIVES

To study surgical and functional outcome of lateral canthal tarsorrhaphy in unilateral paralytic lagophthalmos.

To study improvement in clinical signs and symptoms of exposure keratitis and lid closure after lateral tarsorrhaphy in patients of paralytic lagophthalmos.

## MATERIAL AND METHODS

It was a hospital based prospective case study. Consecutive case series of 10 eyes of 10 patients from Jan. 2012 to Mar. 2013. 10 eyes with unilateral paralytic lagophthalmos with exposure keratitis that didn't respond to conservative medical line of treatment were treated with lateral canthal tarsorrhaphy. Patients having facial nerve paresis with no significant ocular surface changes, non-paralytic lagophthalmos were excluded from the study. Written informed consent was obtained from all patients. Patients were in the age group between 30 to 55 years with mean age of 43.7 years. Most of the patients were males with male to female ratio of 3:2.

Gender	No. of patients
Male	6
Female	4
<b>Total</b>	<b>10</b>

Most of the patients had right sided facial nerve paralysis.

Laterality	No. of patients
Right sided facial nerve palsy	7
Left sided facial nerve palsy	3
<b>Total</b>	<b>10</b>

**CAUSES OF FACIAL NERVE PALSY**

CAUSES	No. of patients
Bell's palsy	7
Trauma (Post Surgical - mastoidectomy)	2
Chronic otitis media	1
Tumors	0
<b>Total</b>	<b>10</b>

**CLINICAL FEATURES**

Patients had facial asymmetry, Loss of forehead wrinkles and nasolabial fold, Drooping of corner of mouth, Difficulty in keeping food in mouth, Drooping of eyebrow, Inability to close the eye, Uncontrolled tearing, Ocular pain and redness in eye. No patients had h/o diabetes.



**Figure 1:** Left sided infranuclear facial nerve palsy with lagophthalmos with exposure of ocular surface



**Figure 2:** Right sided infranuclear facial nerve palsy with lagophthalmos with severe exposure keratitis



**Figure 3:** Left sided facial nerve palsy with lagophthalmos with exposure keratitis

**PRE OPERATIVE ASSESSMENT**

Scleral show especially inferiorly was noted. Pre-operatively it was ranging from 2 – 4mm. Duration of signs and symptoms of exposure keratitis was noted.

**MANAGEMENT**

10 eyes of 10 patients having paralytic lagophthalmos underwent lateral tarsorrhaphy. Indication of tarsorrhaphy

in all patients was to prevent or to treat exposure keratitis. All patients underwent surgery by a single surgeon at GMCH, Nagpur. The surgical steps included marking of intended lateral tarsorrhaphy, splitting of grey line and excision of anterior lamella and suturing of upper and lower limbs of lateral canthal tendon and canthal angle reformation.



**Figure 4:** Post operative photographs

**RESULTS**

10 eyes of 10 patients having paralytic lagophthalmos underwent lateral tarsorrhaphy. Patients were in the age group between 30 to 55 years with mean age of 43.7 years. Most of the patients were males with male to female ratio of 3:2. Laterality - right sided facial nerve palsy was seen in 7 patients and left sided 3 patients. In all cases type of tarsorrhaphy was permanent. 9 patients

needed single tarsorrhaphy and 1 patient needed repeat tarsorrhaphy because of partial opening of tarsorrhaphy. Clinical improvement in signs and symptoms of exposure keratitis was observed in nearly all the cases. Duration of follow-up: 3 months. Mean time to epithelial healing after tarsorrhaphy was 15 days (with range of 10 – 20 days) Inferior scleral show was reduced from 3 mm pre-operatively (range 2 to 4 mm to 0.3 mm post op. with a range of 0 – 0.5 mm at 3 months followup.

## COMPLICATIONS

Recurrence of exposure keratitis was seen in none of the patients. Single patient required repeat tarsorrhaphy all patients were cosmetically satisfied.

## DISCUSSION

The main cause of lagophthalmos is facial nerve paralysis (paralytic lagophthalmos) but it also occurs after trauma or surgery (cicatricial lagophthalmos) or during sleep (nocturnal lagophthalmos). The inability to blink and effectively close the eyes may lead to corneal exposure and excessive evaporation of tear film. While doing tarsorrhaphy it is equally important for the patient to regain a cosmetically acceptable appearance. Conservative treatment includes copious lubrication by eye drops and eye ointment. Other options available are Taping of eyelids, Bandage soft contact lenses, Moisture chamber glasses, Conjunctival flaps, Amniotic membrane grafts, Tissue Adhesives, Glued on hard contact lenses, Blow out patch grafts, Gold weight implants, Palpebral spring insertion, Penetrating keratoplasty, Chemodenervation to yield protective ptosis and Temporalis transplant.

## CONCLUSIONS

Lateral tarsorrhaphy is safe and effective surgical procedure for poor patients having infranuclear Bell's

palsy with severe exposure keratitis, It is simple and effective in achieving functional eye closure providing symptomatic relief and faster corneal epithelial defect healing. Therefore, it is of benefit in cases of patients having exposure keratitis with good success rate and minor complications. Early treatment intervention by lateral tarsorrhaphy is important to help avoid severe complications of exposure keratitis like corneal perforation and endophthalmitis.

## REFERENCES

1. Tan ST et al, Gold weight implantation and lateral tarsorrhaphy for upper eyelid paralysis. J. Craniomaxillofac. Surg. 2013; Apr. 41(3): e49-53.
2. Panda A et al, Lateral Tarsorrhaphy – is it preferable to patching? Cornea 1999 May; 18(3): 299 – 301.
3. Rosenthal P Cotter JM et al, Treatment of persistent corneal epithelial defect with extended wear of a fluid ventilated gas permeable scleral contact lence. AMJ. Ophthalmol. 2000 Jul 130(1): 33 – 41.
4. Cosar CB et al, Tarsorrhaphy clinical experience from a cornea practice. Cornea 2001 Nov; 20(8) 787 – 91.
5. Pushker N Dada et al, Neurotrophic Keratopathy. CLAO J 2001 April; 27(2); 100 – 7.
6. De Silve DJ et al, Surgical technique: modified lateral tarsorrhaphy. Ophthal plast. Reconstr. Surg. 2001 May – June 27 (3) 216 – 8.
7. Gire et al, PROSE treat for lagoph. And expo. Keratopathy ophth. Plast reconst. Surg. 2013 Mar – Apr: 29(2).

Source of Support: None Declared  
Conflict of Interest: None Declared