

Estimation of Height (Stature) From Inferior Extremity Length and Foot Length in Children.

Swati Ramakant Pandhare¹, Anjali D. Patil², A. Kasote³, M. M. Meshram⁴

¹Assistant Professor, Govt. Medical college, Nagpur (MS) INDIA.

²Associate Professor, B.J. Medical College, Pune (MS) INDIA.

³Associate Professor, Govt. Medical College, Nagpur (MS) INDIA.

⁴Professor, Govt. Medical College, Nagpur (MS) INDIA.

Corresponding Addresses:

Swatipandhare84@gmail.com

Research Article

Abstract: Personal identification is an integral part of the investigation in cases of mass disasters where disintegrated and amputated body organs are found very frequently. Estimating stature from various parameters based on the above mentioned evidences becomes one of the most important and essential exercise for personal identification. Different anthropometric parameters have been measured of 1120 children (576 male children, 544 female children) of the age group ranging from Birth to 5 years. Among these findings, height, inferior extremity length and foot length have been evaluated for sexual dimorphism as well as subjected to statistics to study the correlation of inferior extremity length and foot length with that of height. There is scarcity of literature suggesting stature estimation in this age group hence, further the regression equations have been drawn from the data collected. It has been observed that stature can be estimated from the inferior extremity length and foot length. The Technical Error of Measurements was within the accepted limits. It is determined that there was significant differences ($p < 0.05$) in stature, inferior extremity length and foot length between sexes. Stature, inferior extremity length and foot length are positively and significantly correlated with each other ($p < 0.01$).

Keywords: stature, foot length, inferior extremity length, children, personal identification

Introduction:

Determination of sex and estimation of stature from incomplete skeleton as well as from decomposing bodies is the burning issue in forensic science. This has mass application in recent era because of mass disasters like bomb blasts, tsunamis, earthquake, plane crash, mass suicide, forest fires.

It is well known that trunks and limbs exhibit consistent ratios among themselves and to total body height. The ratios between body segments are age, sex and race dependent (Meadows, 1996[1]; Meadows & Jantz, 1999[2]; Williams et al., 2000[3].

This relationship of different body parts is used to establish the sex as well as stature of an individual (Ebite *et al.*, 2008)[4]. This is also very useful anthropologically to find the racial differences, and

medico legally, when only parts of the deceased body are available. Estimation of height of an individual has obvious significance in the personal identification. As height is fundamental to assessing growth and nutrition, calculating body surface area, and predicting pulmonary function in childhood (Gauld et al., 2004)[5].

However, a number of common disabilities and disease processes make it difficult to accurately measure standing height in many patients (Auyeung et al., 2009)[6].

The majority of old methods of estimating stature are limited to measuring whole limb bone and correlating living stature and limb bone length. But few studies are reported in which an attempt has been made to estimate stature from fragmentary or mutilated parts of the body.

Studies on the estimation of stature from the skeletal remains or from the mutilated limbs, mostly of the long bones have been reported as indicated by the published work of the Pearson (1899)[7], Trotter and Glessner (1952)[8]. The Indian perspective of the problem of stature estimation has been studied by the Athwale et al (1963)[9], Patel et al (1964)[10], Joshi et al (1964,65)[11], Lal and Lala (1972)[12], Kalte and Bansal (1974)[13], Thakur and Rai (1987)[14], Saxena (1984)[15], Bhatnagar et al (1984)[16], Jasuja (1987)[17].

Also there are few studies in which stature is estimated from inferior extremity length and foot length. (Agnihotri A.K. et al(2007)[18], Ozden H et al (2005)[19], Sangli S.G. et al (2005)[20],

Oommen A. et al (2005)[21], Krishnan K.(2007)[22], Patel et al (2007)[23].

Among the literature which has searched, there is scarcity of such literature availability in relation to pediatric age group. And hence this information will be highly important to Forensic scientists, human biologists and physical anthropologists for determination of stature from the fragmentary remains of lower limbs of children.

Material and Methods:

1120 children (576 male children, 544 female children) of the age group ranging from Birth to 5 years were examined for different anthropometric norms with informed consent.

This work was carried out at the well baby clinic of the Department of Pediatrics and immunization OPD of department of PSM at Medical College Hospital, and at different primary schools of Municipal Corporation”. Studies on the new born were carried out in the maternity wards of the Medical College Hospital.

The information regarding the exact date of birth was obtained from the hospital and school records and also from the records available with the parents. All measurements were done personally. Those born premature, congenital malformations or diseased children and newborn having birth weight less than 2000 gm were excluded from the study. As the study was carried on the city population which lacks in socio-economic homogeneity, the analysis did not take into account the socio-economic status. Only apparently normal, healthy infants and children were selected for the study.

1) **Height:**

Recumbent length (Crown-heel) was recorded for babies up to 2 years.

The child was placed supine on the “Fiber-glass osteometric board”; the head firmly kept in position against the upright fixed head board; the legs straightened, turning the feet to right angles with the legs keeping the toes upwards. The free foot board was brought into firm contact with the child’s foot and the reading was taken to fraction of centimeter, from the scale set into the measuring table.

Standing height was measured in cases of children from 2 years to 5 years.

The subject is made to stand erect and barefooted on a level ground or platform with the heels together and arms hanging. The occiput, shoulders, buttocks and heels are in the same plane and perpendicular to the ground. The head is so held that the eyes are directed on the horizon. The anthropometric rod is held either in front or back of the subject ensuring that it is perfectly vertical and parallel to the mid-sagittal plane. The cross-arm of the instrument is gently brought down to touch the vertex. Measurements were recorded to the nearest of 0.1cm.

2) **Length of inferior extremity:** This distance was measured from the tip of anterior superior iliac spine to the medial border of sole vertically below the tip of medial malleolus.

3) **Foot length:** It was measured with osteometric board. The distance between prominent points of heel to the tip of great toe was recorded.

Results:

Table I shows, the mean height (*HT*), inferior extremity length (*IEL*) and foot length (*FL*) of the study subjects were significantly different between the genders. Statistical analysis indicated that variation was significant for the measurements of HT, IEL, FL in both sexes. Z value more than 1.96 is statistically significant.

Table I: shows the mean height (Ht), mean inferior extremity length (IEL) and Foot length (FL) of male and female children in the age group of at birth to 5 years.

Age	Mean Height			IEL			FL		
	Male	Female	Z value	Male	Female	Z value	Male	Female	Z value
At birth	48.41	45.8	8.37	21.37	29.65	20.14	7.75	7.48	18.55
Upto 3mths	60.86	57	13.40	24.43	23.01	4.45	8.84	8.44	8.69
3-6 mths	62.25	63	2.54	28.11	27.08	4.24	10	9.43	4.24
6-9 mths	67.23	65.16	7.02	30.82	29.94	5.60	10.82	10.24	5.6
9-12 mths	70.37	69.09	4.33	32.67	31.32	4.93	11.3	10.8	5.21
1yr +	74.00	72.79	8.70	35.98	36.44	3.35	12.38	12.04	10.69
2yr+	80.60	80.32	2.07	42.24	41.06	10.48	14.08	13.81	10.48
3yr+	88.60	87.76	6.56	48.7	47.07	9.65	15.5	14.99	9.65
4yr+	95.13	94.18	6.68	53.47	52.47	9.34	16.27	15.87	9.34




Figure I: shows the mean height (Ht), of male and female children in the age group of at birth to 5 years.




Figure II: shows the mean inferior extremity length (IEL) of male and female children in the age group of at birth to 5 years.




Figure III: shows the mean Foot length (FL) of male and female children in the age group of at birth to 5 years.

Table II: shows the correlation coefficient between height and inferior extremity length and foot length is positive, suggesting that it is significant.

	Height			
	Male		Female	
	Cor. Coe.	P Value	Cor. Coe.	P value
IEL	0.990	0.00	0.937	0.00
FL	0.968	0.00	0.950	0.00

Pearson’s correlation coefficient was used to examine the relationship between, height and inferior extremity length and that of foot length in male and in female children. Correlation coefficient among them was found to be statistically significant $P=0.00$ and positive in both males and females (Table II). It means there is a strong bond between height and inferior extremity length, also between height and foot length and if either of the measurement is known, the other can

be calculated and this would be useful for Anthropologists and Forensic Medicine experts. And hence the regression equations were calculated for male and female.

Table III: shows the **Regression equations** for male and female child:

For Male:

Equation	SE of coefficient	P value
$Ht = 26.5 + 1.29 IEL$	0.00856	0.00
$Ht = 15.1 + 4.81 FL$	0.05870	0.00

For Female:

Equation	SE of coefficient	P value
$Ht = 19.9 + 1.42 IFL$	0.02516	0.00
$Ht = 17.1 + 4.58 FL$	0.07188	0.00

Discussion:

The estimation of height from various long bones, head length and hand length has been attempted by many workers. Singh and Sohal (1951)[24], Jit and Singh (1956)[25], have shown a significant correlation between height and length of clavicle. Athawale (1963)[9] derived a regression equation between total height and forearm bones. Patel, Joshi and Dongre (1964)[11], have derived regression equation between tibia and total height in Gujarati population.. Saxena et al (1981)[15], derived a regression equation between head length and height and find significant correlation coefficient, + 0.2048.

Regarding the estimation of height from inferior extremity length and foot length, Sultan et al (2005)[26] was carried out study to estimate the relationship between hand length, foot length and stature using multiple linear regression analyses. Qamra et al (1979)[27], derived regression equation between foot length and height in northwest India od adult age group. Also Patel et al (2007)[10], Chikhalkar B.G.[28], given similar regression equation from foot length.

However lower limb dimensions of pediatric age group have not frequently been used for this. The present study deals with the observations on correlation of total standing height with inferior extremity length and foot length of pediatric age group from at birth to 5yrs of age.

The average height of adult males within a population is significantly higher than that of adult females (Williams *et al.*)[3]. The results obtained in this study are in agreement with the above statement. There was distinct sexual dimorphism in the height, inferior extremity length and foot length

in our study group also. The need for gender specific formulae is proved as the rate of skeletal maturity in males and females tend to vary during the course of development. The regression analysis proved to be the easiest and the reliable method (Meadows)[2].

Regression models for stature prediction were formulated using height, inferior extremity length and foot length lengths and checked for their accuracy by comparing the estimated stature and the actual stature.

As variety of factors such as, age, race, gender and nutritional status affect human development and growth and therefore, different nomograms are required for different populations (Williams *et al.*; Joshi *et al.*). The present study for the first time documents norms for height with inferior extremity length and foot length and presents gender specific linear regression models for stature prediction in pediatric population. The formulae are valid for the age group from birth upto 5 yrs of age of the subjects.

The regression models proposed will be of immense practical use in clinical practice, medico-legal, anthropological and archeological studies where the total height of a subject can be calculated if the superior extremity length or hand length is known.

Limitations:

1. In the present study, age range of only at birth to 5 years is considered.
2. Measurements of only healthy children are considered. Hence the data may not be applicable to children who are malnourished &/or suffering from congenital structural malformations.
3. The present study is a preliminary one & would be followed up by other similar studies to address the above limitations.

Acknowledgment:

My sincere thanks to head of department, senior teachers, colleagues of department of anatomy, B. J. Medical college, Pune, and Govt. Medical college, Nagpur. Maharashtra, India. Also my sincere thanks to Statistician, Dept PSM, Govt. Medical College, Nagpur.

Declaration for Ethical standards:

Dr Swati Ramakant Pandhare, hereby declare that during the work of present study precautions are taken for not to violate all laws of India to maintain ethical standards.

Conflict of interest:

The authors declare that they have no conflict of interest.

References:

- [1] Meadows, L. Secular change and allometry in the long limb bones of Americans from the mid 1700 through the 1970s. Knoxville, Tennessee, University of Tennessee, 1996.
- [2] Meadows, L. & Jantz, R. L. Secular changes in long bone length and proportion in the United States 1800-1970. *Am. J. Phys. Anthropol.*, 110:57-67, 1999.
- [3] Williams, P. L.; Bannister, L. H.; Berry, M. M.; Collins, P.; Dyson, M. & Dussek, J. E. *Gray's Anatomy: The Anatomical basis of medicine and surgery*. 38th Ed. New York, Churchill Livingstone, 2000.
- [4] Ebite, L. E.; Ozoko, T. C.; Eweka, A. O.; Otuaga, P. O.; Oni, A. O. & Om'Iniabohs, F.A.E. Height: Ulna Ratio: A Method of Stature Estimation In A Rural Community in Edo State, Nigeria. *The Internet Journal of Forensic Science*, 3(1), 2008.
- [5] Gauld, L. M.; Kappers, J.; Carlin, J. B. & Robertson, C.F. Height prediction from ulna length. *Dev. Med. Child Neurol.*, 46(7):475-80, 2004.
- [6] Auyeung, T. W.; Lee, J. S. W.; Kwok, T.; Leung, J. Leung, P.C. & Woo, J. Estimation of stature by measuring fibula and ulna bone length in 2443 adults: *J. Nut. Health Aging.*, 10:931-6, 2009.
- [7] Pearson K. Mathematical contributions to the theory of evolution V. On the reconstruction of stature of pre historic races. *Philosophical Transactions of Royal Society; London; 1899; 192: 169-224.* Cited in "Krogman's Human Skeleton in Forensic Medicine" 2nd Edition, Charles Thomas, U.S.A., 1899.
- [8] Trotter M and Glesser GC. Estimation of stature from long bones of American White and Negroes, *American journal of physical anthropology* 1952; 10: 463, 1952.
- [9] Athwale MC. Estimation of height from length of forearm bones - A study on 100 Maharashtrian male adults of age 25-30, *American journal of physical anthropology*, 1963; 21: 105-112, 1963.
- [10] Patel MP, Joshi NB and Dongre AV. Regression equation of height on tibial length. *Indian Journal of Medical Research*, 1964; 52: 531-534, 1964.
- [11] Joshi, N. B.; Patel, M. P. & Dongre, A. V. Regression equation of height from ulna length. *Ind. J. Med. Res.*, 52:1088-91, 1964.
- [12] Lal CS and Lala JK. Estimation of stature from tibial and ulnar lengths in North Bihar. *Journal of Medical Association; 1972; 58: 120-121, 1972.*
- [13] Kalte PM and Bansal PC. Determination of regression formulae for reconstruction of stature from long bones of upper limbs in Maharashtrians and Marathwara region. *Journal of Anatomical Society (India)*, 1974; 23: 6-11, 1974.
- [14] Thakur SD and Rai KS. Determination of Stature from hand measurement, *Medicine Science and Law*, 1987; 78: 25-28, 1987.
- [15] Saxena SK. Study of correlations and estimation of stature from hand length, hand breadth and sole length. *Anthropol Anz.* 1984; 42(4):271-6, 1984.
- [16] Bhatnagar DP, Thapar SP and Batish MK. Identification of personal height from somatometry of hands in the Punjabi males, *Forensic science international*, 1984; 24 : 137- 141, 1984.
- [17] Jasuja OP. Calculation of stature from foot and shoe impressions, PhD Thesis, Punjabi University, Patiala. (unpublished) 1987.
- [18] Agnihotri AK, Shukla S, Purwar B. Determination of sex from the foot measurements. *The Internet J Forensic Sci.*, 2:1, 2007.
- [19] Ozden H, Balci Y, Demirustu C, Turgut A, Ertugrul M. Stature and sex estimate using foot and shoe dimensions. *Forensic Sci Int.*, 147:181-4, 2005.
- [20] Sanli SG, Kizilkanat ED, Boyan Ozsahin NE, M. Bozkir MG, Soames R, Erol H, and Oguz O. Stature Estimation Based on Hand Length and Foot Length. *Clin Anat.*, 18:589-596, 2005.
- [21] Oommen A, Mainker A, Oommen T. A Study of the Correlation Between Hand Length And Foot Length In Humans. *J Anat Soc. India*, 54:1-9, 2005.
- [22] Krishan K, Sharma A. Estimation of stature from dimensions of hands and feet in a North Indian population. *J Forensic Legal Med.*, 14:327-332, 2007.
- [23] Patel S. M., Shah G. V., Patel S. V. Estimation of height from measurement of foot length in Gujrat region. *J Anat Soc India*, 56(1): 25-27, 2007.
- [24] Singh B and Sohal HS. Estimation of stature from clavicle in Punjabis. A preliminary report. *Indian Journal of Medical Research*, 40: 67 – 71, 1951.
- [25] Jit Inder and Shamer Singh. Estimation of stature from clavicle. *Indian Journal of Medical Research*, 44: 137-155, 1956.
- [26] Sanli SG, et. al. Stature estimation based on hand length and foot length. *Clinical Anatomy*, 18; 8: 589–596, 2005.
- [27] Suneel Qamra, Inderjit and Deodhar SD. A model for reconstruction of height from foot measurements in a adult population of Northwest India. *Indian Journal of Medical Research*, 71: 77-83, 1980.
- [28] Chikhalkar B.G., Mangaonkar A.A., Nanandkar S.D., Peddewad R.G. Estimation of Stature from Measurements of Long Bones, Hand and Foot Dimensions. *J Indian Acad Forensic Med*, 32(4): 329-33, 2010.