

Glucose and Total Sugars Content and Changes therein in the Seeds of *Schleichera oleosa* (Lour.) Oken.(kusum) due to Bioteriation by Pathogenic Fungi during Storage

AK Srivastava* and GK Pandey**

*Head, Dept of Botany, St .Xavier's College, Ranchi INDIA

**Dept of Botany, St Columba's PG College, Hazaribag, INDIA

Corresponding Address:

ajaysrivastava11@gmail.com

Research Article

Abstract: Seeds of one of the non-edible oilseed, kusum (*Schleichera oleosa* f. Sapindaceae) are of great significance in Jharkhand area and its oil is used in burning lamps, varnishing, massage and medicine. Its oil-cake is a good manure. The unsanitary and humid condition makes it prone to a faster biodeterioration due to several fungal inhabitants during storage. The colonising fungi in the seeds use up some of the glucose and sugars as carbon sources for meeting energy requirements in their prolonged association with the seeds. In the span of one year the glucose level dropped by as much as 50.2 percent of its initial value. Similarly, total sugar also registered a loss of 40.4 percent due to the association of the mycodwellers.

Keywords: *Schleichera oleosa*, oilseed, colonising fungi, carbon sources, mycodwellers.

Introduction:

Seed pathology is up and coming area of plant pathology and has attained a great magnitude. Seeds of a non edible oil, *Schleichera oleosa* f. Sapindaceae (Kusum) are of great use in burning lamps, varnishing, massage and medicine. Its oil-cake is good manure. In the Jharkhand area, Kusum seeds are collected just after onset of rainy season i.e. by the end of month of May. Conventionally seeds are stored in gunny bags in the village houses and godowns. In either case the condition is unsanitary and humid and is conducive for fungal growth leading to good scale destruction.

There are many reports indicating that fungi are associated with seed surface as well as internal mycoflora (DUTTA & ROY , 1987 ; JAYWEERA *et*

al 1988). It was worthwhile to seek the extent of biodeterioration occurring in the seeds in terms of soluble carbohydrates in different forms viz. glucose and total sugars.

Material and Methods:

Kusum seeds were obtained from the godown of Ranchi forest department for the preparation of oil and oilcake on regular interval of one month. The work spanned from June 2001, one harvest season to May 2002, just before the other harvest season. Glucose content was estimated by titration method using Fehling solutions (Cole's method, 1933). Total sugars was estimated by colorimetric method using anthrone as a reagent (DUBOIS *et. al.*, 1951) using a colorimeter of Systronics (India) make. Estimations were done in triplicate and standard deviation and standard errors were calculated.

Results and Discussion:

The effect of continued fungal activity on the seeds manifested in the lowered glucose, total soluble sugars and starch contents just like oils (SRIVASTAVA ,AK and PANDEY GK. 2000). The glucose level dropped by as much as 50.2 percent in the span of one year of its initial value. Similarly, total sugar also registered a loss of 40.4 percent due to the association of the mycodwellers. The results are elaborated in figs 1-2.

Table 1: Amount of glucose present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack , June '01- May '02.

S No	Month	Wt.(mg)	Percentage	Change(mg)
1	Jun '01	16.75	1.675	-----
2	Jul '01	16.25	1.625	-0.05
3	Aug '01	16.25	1.625	-0.05
4	Sept '01	15.75	1.575	-0.1
5	Oct '01	14.9	1.49	-0.175
6	Nov '01	14.3	1.43	-0.245
7	Dec '01	13.1	1.31	-0.365
8	Jan '02	13.0	1.3	-0.375
9	Feb '02	11.9	1.19	-0.485
10	Mar '02	10.3	1.03	-0.645
11	Apr '02	9.5	0.95	-0.725
12	May '02	8.35	0.835	-0.84
	SD: 0.284	SE: 0.081		

Final percent change in glucose content $[(1.675 - 0.835)/1.675] 100 = 50.2\%$




Fig 1: Amount of glucose present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack , June '01- May '02.

Table 2: Amount of sugars present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack , June '01 - May '02

S No	Month	Wt.(mg)	Percentage	Change(mg)
1	Jun '01	47	4.7	-----
2	Jul '01	47	4.7	-0
3	Aug '01	45	4.5	-0.2
4	Sept '01	44	4.4	-0.3
5	Oct '01	41	4.1	-0.6
6	Nov '01	40	4.0	-0.7
7	Dec '01	37	3.7	-1.0
8	Jan '02	35	3.5	-1.2
9	Feb '02	34	3.4	-1.3
10	Mar '02	32	3.2	-1.5
11	Apr '02	30	3.0	-1.7
12	May '02	28	2.8	-1.9
	SD: 0.663	SE: 0.191		

Final percent change in sugars content $[(47-28)/47]100 = 40.425\%$




Fig 2: Amount of sugars present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack, June '01 - May '02

Changes in glucose content with pathogenic association have been reported. CHAUDHARY and PRASAD (1974) found a depletion of glucose among other nutrients in the shoot tissues of pigeon pea infected with *Fusarium oxysporum* f.sp. *udum*. CHATTOPADHYAYA and NANDY (1978) a carbohydrate loss to *F. subglutinum* infection. Loss in amount of glucose in fruits have been reported for tomato- *Drechslera australiense* (KAPOOR and TANDON, 1970); tomato- *Alternaria solani* (MEHTA et. al., 1975); banana- *Gloeosporium musarum* (WANG, 1960) and citrus- *Xanthomonas campestris* pv. *Citri* (VIDYASEKARAN and DURAIRAJ, 1971) host pathogen systems.

Changes in total sugar content in the host tissues due to pathogenic actions have been reported by CRAIG and HOOKER (1961); DHANVANTARI (1967); DAYAL and JOSHI (1968) and PADMANABHAN (1988) on different host pathogen systems.

The drop is due to the sugar being used by the fungal pathogens as respiratory substrates (BAKER, 1965; WU, 1973). All these reports are concurrent to our findings. It is thus apparent that one year of seed infestation of *Schleichera oleosa* predominantly and jointly by *Aspergillus fumigatus*, *A. flavus*, *A. niger*, *Fusarium solanii*, *Paecilomyces variotii* and *Mucor sp.* causes biodeterioration of its seeds manifested in the diminishing of soluble carbohydrate contents.

References:

- [1] Chattopadhyay NC., and Nandy B. 1978. Changes in nitrogen content in malformed inflorescence of mango caused by *Fusarium moniliforme* Sheld. Var *subglutinans* Wr. Et. Rg *Giornale Bot. Ital.* 112: 343-346.
- [2] Cole, SW. 1919. The estimation of lactose and glucose by the copper *Biochem J.* 13(2).
- [3] Craig J and Hooker AL. 1961. Relation of sugar trends and pith density to *Diplodia* stalk rot in dent corn. *Phytopathology* 51 : 376- 382
- [4] Dayal R and Joshi MM. 1968. Post infection changes in the sugar content of leaf spot infected barley. *Indian Phytopath.* 21 : 221-222.
- [5] Dubois, M., Gilles, K., Hamilton, J.K., Rebers, PA and Smith, F., 1951 A colorimetric method for the estimation of sugar. *Nature* 168: 167.
- [6] Dhanvantri, BN. 1967. The leaf scorch disease of Strawberry, *Diplocarpon earlianum* and the nature of resistance to it. *Can. J. Bot.* 45 : 1525-1543
- [7] Kapoor, IJ and Tandon RN. 1970. Post infection changes in sugar content of tomato fruits caused by *Drechslera australiense*. *Indian Phytopath.* 23 : 133-135.
- [8] Mehta P., Vyas, KM and Saksena, SP. 1975. Metabolic changes during pathogenesis of fruit rot disease of tomato. *Indian Phytopath.* 28 : 253- 255.
- [9] Padmanabhan, P., Alexander, KC. and Shanmugam, N. 1988. Some metabolic changes induced in sugarcane by *Ustilago scitaminea*. *Indian Phytopath.* 41: 229-232.
- [10] Srivastava, AK and Pandey, GK. 2000. Chemical changes in properties of kusum (*Schleichera oleosa*) oil during its seed infestation by fungi. *J. Mycopathol. Res.* 38(1): 29-32, (2000)
- [11] Wang, MC 1960. Physiological studies on *Gloeosporium musarum* Cook. At Mass, the causal organism of banana anthracnose. Changes in the carbohydrate composition of banana pulp with reference to the adaptive secretion of *amylase*. *Bot. Bull. Acad. Sin. N.S.* 1 : 59- 75.