

A Study of clinical profile of the patients with myocardial ischemia at tertiary health care centre

Pankaj Bohra^{1*}, Eshan Sharma², Kamal Kumar Jain³

{¹Assistant Professor, Department of Cardiology} {^{2,3}Assistant Professor, Department of Medicine}
NIMS Medical College an Hospital, Jaipur, Rajasthan, INDIA.

Email: jaindrkamalkumar@gmail.com

Abstract

Introduction: In 2003, the prevalence of CHD in India was estimated to be 3-4 per cent in rural areas (two-fold higher compared with 40 year ago), and 8-10 per cent in urban areas (six fold higher compared with 40 year ago), with a total of 29.8 million affected. **Aims and Objectives:** To study clinical profile of the patients with myocardial ischemia at tertiary health care Centre. **Methodology:** This was a hospital based cross-sectional study of the Patients who were admitted to tertiary health care Centre with diagnosis of Myocardial Ischemia diagnosed by ST-Elevation Myocardial Infarction during the Period of One year from June 2014 to June 2021 at tertiary health care Centre. Total 110 patients were enrolled into the study. All the Socio-demographic information and Clinical information were recorded and the associated Risk factors have also studied. **Result:** The majority of the Patients were from 40-50- 25.45 % followed by 50-60- 23.63 %; 30-40- 20.90 %; >60- 19.09 %; 20-30- 10.90 % Majority of the Patients were Male i.e. 56.36% and 43.63% were Female Most common associated factors with MI patients were H/o Smoking - 52.72% followed by H/o Hypertension -48.18%; Obesity (BMI >30) -39.09%; H/o Tobacco Chewing -34.45%; H/o Diabetes - 22.72%; Hypercholesteremia -21.81% Out of the Total 110 Patients 82.72%- Recovered; 10.90%- Discharge Against medical Advise; 6.36%- Died. **Conclusion:** The most common age of presentation of Myocardial infarction observed in our study is 40-50 and was associated with risk factors like Smoking, Hypertension, Obesity, Tobacco Chewing, Diabetes, Hypercholesteremia.

Key Words: Myocardial ischemia, Obesity, Diabetes, Tobacco Chewing, CHD(Coronary Heart Disease).

*Address for Correspondence:

Dr. Pankaj Bohra, Assistant Professor, Department of Cardiology, NIMS Medical College an Hospital, Jaipur, Rajasthan, INDIA.

Email: jaindrkamalkumar@gmail.com

Received Date: 02/02/2021 Revised Date: 21/03/2021 Accepted Date: 30/04/2021

Access this article online

| | |
|---|--|
| Quick Response Code: | Website: www.statperson.com |
|  | Volume 11 Issue 2 |

INTRODUCTION

In 2003, the prevalence of CHD in India was estimated to be 3-4 per cent in rural areas (two-fold higher compared with 40 year ago), and 8-10 per cent in urban areas (six fold higher compared with 40 year ago), with a total of 29.8 million affected (14.1 million in urban areas, and 15.7 million in rural areas) according to population-based cross-sectional surveys.^{1,2} The Global Burden of Diseases

(GBD) study reported the estimated mortality from coronary heart disease (CHD) in India at 1.6 million in the year 2000.³ A total of nearly 64 million cases of CVD are likely in the year 2021, of which nearly 61 million would be CHD cases (the remaining would include stroke, rheumatic heart disease and congenital heart diseases). Deaths from this group of diseases are likely to amount to be a staggering^{3,4} million.⁴ Driving this steep rise in CVD risk factor burden is the rapid increase in the proportion of urban inhabitants (currently at 30% with a projected rise to 43% in 2021).⁵ Urbanization is characterized by a marked increase in the intake of energy-dense foods, a decrease in physical activity, and a heightened level of psychosocial stress, all of which promote the development of dysglycaemia, hypertension, and dyslipidaemia.⁶

METHODOLOGY

This was a hospital based cross-sectional study of the Patients who were admitted to tertiary health care Centre with diagnosis of Myocardial Ischemia diagnosed by ST-

Elevation Myocardial Infarction during the Period of One year from June 2014 to June 2021 at tertiary health care Centre. Total 110 patients were enrolled into the study. All the Socio-demographic information and Clinical information were recorded and the associated Risk factors have also studied.

RESULT

Table 1: Age wise Distribution of the MI Patients

| Age | No. | Percentage (%) |
|--------------|------------|-----------------|
| 20-30 | 12 | 10.90 % |
| 30-40 | 23 | 20.90 % |
| 40-50 | 28 | 25.45 % |
| 50-60 | 26 | 23.63 % |
| >60 | 21 | 19.09 % |
| Total | 110 | 100.00 % |

The majority of the Patients were from 40-50- 25.45 % followed by 50-60- 23.63 %; 30-40- 20.90 %; >60- 19.09 %; 20-30- 10.90 %

Table 2: Gender wise Distribution of the MI Patients

| Sex | No. | Percentage (%) |
|--------------|------------|----------------|
| Male | 62 | 56.36% |
| Female | 48 | 43.63% |
| Total | 110 | 100% |

Majority of the Patients were Male i.e. 56.36% and 43.63% were Female

Table 3: Distribution of the Patients as per the Associated Risk Factors

| Risk Factors | No. | Percentage (%) |
|---------------------|-----|----------------|
| H/o Smoking | 58 | 52.72% |
| Obesity (BMI >30) | 43 | 39.09% |
| Hyper-cholestremia | 24 | 21.81% |
| H/o Tobacco Chewing | 38 | 34.45% |
| H/o Diabetes | 25 | 22.72% |
| H/o Hypertension | 53 | 48.18% |

Most common associated factors with MI patients were H/o Smoking - 52.72% followed by H/o- Hypertension - 48.18%; Obesity (BMI >30) -39.09%; H/o Tobacco Chewing -34.45%; H/o Diabetes - 22.72%; Hyper-cholestremia -21.81%

Table 4: Distribution of the Patients as per the Outcome

| Outcome | No. | Percentage (%) |
|----------------------------------|------------|----------------|
| Recovered | 91 | 82.72% |
| Died | 7 | 6.36% |
| Discharge Against medical Advise | 12 | 10.90% |
| Total | 110 | 100% |

Out of the Total 110 Patients 82.72%- Recovered; 10.90%- Discharge Against medical Advise; 6.36%-Died.

DISCUSSION

The burden of CVD and its risk factors in India calls for a sound public health approach to stem the epidemic. Efforts to put in place an intervention programme should

be complemented with a robust surveillance mechanism so as to monitor evaluate and guide policies and programmes⁷. The elderly with acute myocardial infarction (AMI) have been reported to present with more atypical symptoms.^{8,9} AMI is associated with significantly higher mortality in the elderly compared with the young,⁹⁻¹⁴ yet the elderly are treated less aggressively than the young.¹⁶ Thrombolytic therapy has the greatest effect in the elderly even though there is an increased risk of haemorrhagicstroke.^{15,16}

The majority of the Patients were from 40-50- 25.45 % followed by 50-60- 23.63 %; 30-40- 20.90 %; >60- 19.09 %; 20-30- 10.90 % Majority of the Patients were Male i.e. 56.36% and 43.63% were Female Most common associated factors with MI patients were H/o Smoking - 52.72% followed by H/o- Hypertension -48.18%; Obesity (BMI >30) -39.09%; H/o Tobacco Chewing -34.45%; H/o Diabetes - 22.72%; Hyper-cholestremia -21.81% Out of the Total 110 Patients 82.72%- Recovered; 10.90%- Discharge Against medical Advise; 6.36%- Died ; death occurred mostly in old aged persons and in whom treatment was delayed and associated with co-morbidity like diabetes, Hypertension, Obesity.

CONCLUSION

The most common age of presentation of Myocardial infarction observed in our study is 40-50 and was associated with risk factors like Smoking, Hypertension, Obesity

Tobacco Chewing ,Diabetes, Hyper-cholestremia.

REFERENCES

1. Gupta R. Burden of coronary heart disease in India. Indian Heart J. 2005;57:632-8.
2. Gupta R. Coronary heart disease in India: Absolute numbers and economic burden. Rapid response to Ghaffar A, Reddy KS, Singhi M. Burden of non-communicable diseases in South Asia.BMJ. 2004;328:807-10.
3. Murray CJL, Lopez AD. Mortality by cause for eight regions of the world: Global burden of disease study. Lancet. 1997;349:1269-76.
4. Ministry of Health and Family Welfare. Burden of disease in India. In: MHFW, eds. Background Papers for the National Commission on Macroeconomics. New Delhi: Ministry of Health and Family Welfare, Government of India; 2005.
5. Reddy KS, Shah B, Varghese C, Ramadoss A. Responding to the threat of chronic diseases in India. Lancet. 2005;366:1746-51.
6. Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases: part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization. Circulation. 2001;104:2746-53.

7. Bela Shah, Prashant Mathur. Surveillance of cardiovascular disease risk factors in India: the need and scope. *Indian J Med Res.* 2010 Nov; 132: 634-42.
8. Bayer AJ, Chadha JS, Farag RR, Pathy MS. Changing presentation of myocardial infarction with increasing old age. *J Am Geriatr Soc.* 1986;34:263e266.
9. Tresch DD, Brady WJ, Aufderheide TP, Lawrence SW, Williams KJ. Comparison of elderly and younger patients with out-of-hospital chest pain. Clinical characteristics, acute myocardial infarction, therapy and outcomes. *Arch Intern Med.* 1996;156:1089e1093.
10. Wang SW, Ren GC, Shio SF, Yu SY, Zhen FY. Acute myocardial infarction in elderly Chinese. A clinical analysis of 631 cases and comparison with 389 younger cases. *Jpn Heart J.* 1998;29:301e307.
11. Day JJ, Bayer AJ, Pathy MS, Chadha JS. Acute myocardial infarction: diagnostic difficulties and outcome in advanced old age. *Age Ageing.* 1987;16:239e243.
12. Yang XS, Willems JL, Pardaens J, De Geest H. Acute myocardial infarction in the very elderly. A comparison with younger age groups. *Acta Cardiol.* 1987; 42:59e68.
13. Paul SD, O'Gara PT, Mahjoub ZA, *et al.* Geriatric patients with acute myocardial infarction: cardiac risk factor profiles, presentation, thrombolysis, coronary interventions and prognosis. *Am Heart J.* 1996;131:710e715.
14. Mehta RH, Rathore SS, Radford MJ, Wang Y, Wang Y, Krumholz HM. Acute myocardial infarction in the elderly: differences by age. *J Am Coll Cardiol.* 2001;38:736e741.
15. Reardon M, Camm JA. CME paper: acute myocardial infarction in the elderly. *Am J Geriatr Cardiol.* 2000;9:138e142.
16. Haase KK, Schiele R, Wagner S, *et al.* In-hospital mortality of elderly patients with acute myocardial infarction: data from the MITRA (Maximal Individual Therapy in Acute Myocardial Infarction) Registry. *Clin Cardiol.* 2000;23:831e836.

Source of Support: None Declared
Conflict of Interest: None Declared