

Effectiveness of silver dressings in treatment of diabetic ulcer

Saurabh Mishra^{1*}, John Joseph S Martis²

¹Post-graduate, ²Professor, Department of General Surgery, Father Muller Medical College, Mangalore, Karnataka, INDIA.

Email: martisjohn@yahoo.co.in, sm9900787060@gmail.com

Abstract

Background and objective: With the increase in the life expectancy, the incidence of diabetic ulcers is inevitable. In this study we compare the efficacy of topical silver preparation with conventional wound dressing in healing of diabetic ulcers, in terms of days required for healing, rate of granulation tissue formation, quality of graft bed, graft uptake, effect on bacterial growth and side effects of silver preparations. **Method:** 58 patients with diabetic ulcers were selected by purposive sampling technique in Father Muller Medical College and Hospital between August 2012 to September 2014. Data related to the objectives of the study were collected. **Result:** this study has shown better granulation tissue formation, graft uptake, negative bacterial growth and decreased hospital stay in patients receiving silver dressings than patients receiving conventional dressings. **Conclusion:** The study concludes that silver preparations are better than conventional preparation. They are also cheap, readily available and easy to use alternative with no side effect.

Keywords: Diabetic ulcer, Silver dressing, Conventional dressings.

*Address for Correspondence:

Dr. Saurabh Mishra, Post-graduate, Department of General Surgery, Father Muller Medical College, Mangalore, Karnataka, INDIA.

Email: sm9900787060@gmail.com

Received Date: 12/05/2021 Revised Date: 20/06/2021 Accepted Date: 26/07/2021

Access this article online	
Quick Response Code:	Website: www.statperson.com
	Volume 11 Issue 3

INTRODUCTION

Wound healing is a complex biological process influenced by several agents, which promote granulation tissue regeneration and contribute towards wound healing. Chronic skin wounds affect approximately 3% of persons elderly and are usually related to neuropathy (eg, diabetic foot or pressure ulcers), vasculopathy (venous stasis or arterial insufficiency ulcers), or trauma. Patients with diabetic ulcers are frequently treated with either systemic or topical antimicrobial therapy. Various studies done have shown that even if the infection remains confined to superficial tissues, not only does it delay wound healing, cause exudation and malodor but in due course of time progresses to involve deeper tissues and may be a potential for systemic infection. Topical therapy

affords many potential advantages like avoidance of systemic adverse effects of the drug. The most simple of all therapeutic procedures which uses the topical antimicrobials is the treatment of cutaneous infection with topical medication. Hence we have decided to study the role of topical silver dressing in healing of Diabetic ulcers.¹ The topical antimicrobial agent silver has been used for hundreds of years in wound care¹ but its use was cumbersome as the silver available for dressings was difficult to apply. Recently of a wide range of wound dressings that contain elemental silver or a silver-releasing compound have been developed which are easier to apply, provide sustained availability of silver, need less frequent dressing changes have overcome some of the problems associated with the first silver preparations². Silver dressings maintain a moist wound environment and help in facilitation of autolytic debridement³.

Diabetic foot ulcer^{7,8,9,10}

It is estimated that in all diabetic 15% of them develop chronic ulcer, and about 25% of those progress to need an amputation. In diabetic ulcers although vascular and neurological impairment is the main reason for reduced healing, recent studies have shown wound healing is impaired in part due to deficiency of growth factors. The treatment of diabetic ulcers requires frequent

debridement, moist wound care, treatment of infection, and revascularization of ischemic limb if necessary. Silver ions are highly reactive and affect multiple sites within bacterial cells, ultimately causing bacterial cell death by binding to the bacterial cell membranes and causing disruption of the bacterial cell wall and cell leakage, they also interfere with energy production, enzyme function and cell replication and thus are active against a broad range of bacteria, fungi and viruses.^{4,5}

OBJECTIVES OF STUDY

- To study the efficacy and advantages of topical application silver in diabetic ulcer.
- Comparison between topical silver preparations and conventional dressings in management of diabetic ulcer.

MATERIALS AND METHODS

Source of Data

Patients in Father Muller Medical College Hospital, Mangalore admitted with diabetic ulcers between august 2012 and September 2014.

Method of Collection of Data

Study type:

Prospective study The study was done on 58 patients who presented with diabetic ulcers at Father Muller hospital with purposive sampling technique. A complete detailed history and physical evaluation, relevant blood investigations, radiological investigations were done.

Data Collection:

Age of the patient, Symptoms and their duration, Past history, Complete physical examination, laboratory investigations (including specific investigations like Pus cultures was done at regular intervals), Radiological investigations wherever necessary was done and treatment also given.

Inclusion Criteria: All patients above 20 yrs who were diagnosed to have diabetic ulcers were included.

Exclusion Criteria: Malignant ulcers, X-rays showing features of osteomyelitis, Other clinically significant medical conditions that would impair wound healing including renal .hepatic, hematological, neurological and immunological diseases, Patients receiving corticosteroids, immunosuppressive agents, radiation, or chemotherapy within one month prior to entry into study were also excluded.

Plan for Data - Analysis

Collected data will be analyzed by ANOVA for repeated measures and Chi - Square test.


RESULTS

In our study 58 patients were chosen and further divided into a study group consisting of 29 patients who were designated to receive the silver dressings and a control

group consisting of 29 patients who were designated to receive the conventional dressings after elimination of all statistically significant bias.

Table 1: Discharge (purulent)


Discharge	Silver	Betadine
Day 0	29	29
Day 5	22	25
Day 10	12	20



As we can see in our study the amount of discharge is decreased more in case of silver dressings as compared to conventional dressings.

Table 2: Slough


Slough	Silver	Betadine
Day 0	29	29
Day 5	20	24
Day 10	13	19



As we can see in our study the quantity of slough is decreased more in case of silver dressings as compared to conventional dressings.

Table 3: Granulation tissue

Granulation tissue	Silver	Betadine
Day 0	0	0
Day 5	18	13
Day 10	25	19



As we can see in our study that granulation tissue formation is more in case of silver dressings as compared to conventional dressings.

DISCUSSION

This study has shown better granulation tissue formation, reduced slough formation, decreased purulent discharge, better graft uptake, and negative bacterial growth and decreased hospital stay in patients receiving silver dressings than patients receiving conventional dressings which is in comparison with Sergensen. B et al;⁶ conducted an open label multicentre randomised controlled study in which they evaluated silver releasing alginates foam dressings with foam dressings in critically colonised chronic ulcers and found that silver releasing alginate foam dressings reduced ulcer area significantly, fewer leakages and lesser maceration was seen. In another study by Eun⁷ and colleagues demonstrated that silver nanoparticles had a broad-spectrum antimicrobial activity and could also be used as a topical agent applied to cervical tissues or to mucous membrane.

CONCLUSION

The study concludes that silver preparations are better than conventional preparation. They are also cheap, readily available and easy to use alternative with no side effect.

REFERENCES

1. Lansdown ABG. A review of the use of silver in wound care: facts and fallacies. *Br J Nurs*. 2004; 13(6): S6-S19.
2. Parsons D, Bowler PG, Myles V, Jones S. Silver antimicrobial dressings in wound management: a

- comparison of antibacterial, physical, and chemical characteristics. *Wounds*. 2005; 17(8): 222-32.
3. Woodward M. Silver dressings in wound healing: what is the evidence? *Primary Intention*. 2005; 13(4): 153-60.
4. Coates AR, Halls G, Hu Y. Novel classes of antibiotics or more of the same. *Br J Pharmacol*. 2011; 163(1): 184-94.
5. Percival SL, Bowler P, Russell D. Bacterial resistance to silver in wound care. *J Hosp Inf*. 2005; 60: 1-7.
6. Jorgensen B, Price P, Andersen KE et al. The silver releasing foam dressing, contact foam promotes faster healing of critically colonized venous leg ulcers, randomized controlled study. *INT wound journal*. 2005; 2(1):64-73.
7. PHYSIOLOGY OF WOUND HEALING From: Habif: Clinical Dermatology, 5th ed.; Chapter 27 - Dermatologic Surgical Procedures. WOUND HEALING.
8. A. Willrich, M. Pinzur, M. McNeil, D. Juknelis, and L. Lavery, "Health related quality of life, cognitive function, and depression in diabetic patients with foot ulcer or amputation. A preliminary study," *Foot and Ankle International*, vol. 26, no. 2, pp. 128-134, 2005
9. M. A. Loot, S. B. Kenter, F. L. Au et al., "Fibroblasts derived from chronic diabetic ulcers differ in their response to stimulation with EGF, IGF-I, bFGF and PDGF-AB compared to controls," *European Journal of Cell Biology*, 2002:81; pp. 153-60
10. J. Apelqvist, G. Ragnarson-Tennvall, U. Persson, and J. Larsson, "Diabetic foot ulcers in a multidisciplinary setting. An economic analysis of primary healing and healing with amputation," *Journal of Internal Medicine*. 1994:235; pp. 463-71.

Source of Support: None Declared
Conflict of Interest: None Declared