

Adult type ureterocele in lower moiety of duplicated draining system- case report

S R Kulkarni¹, Ashar Shaikh^{2*}, Ronak Chaudhari³, Sandeep Gupta⁴, Tejas Chincholi⁵, Kishor Bandagar⁶

¹Professor, ^{2,3,4,5,6}Resident, Department of General Surgery, Krishna Institute of Medical Sciences, Karad, Maharashtra, INDIA.
Email: ashars97@gmail.com

Abstract

Ureterocele is dilatation of lower ureter at its entry into urinary bladder. It is classified as single or duplex system and acquired or congenital. Acquired type is commonly seen in adults. Duplicated ureters are often associated with an obstructed upper pole moiety and a refluxing lower pole moiety. In our case report adult type simple ureterocele is presented in lower moiety of duplex system with impacted calculus. Because of rarity of the presentation and diagnosis of the condition, urologist needs to be aware of etiology and appearance. Treatment options depend on the extension of the ureterocele, obstruction to draining system.

Key words: ureterocele, duplex system, adult type ureterocele.

*Address for Correspondence:

Dr. Ashar Shaikh, Resident, Department of General Surgery, Krishna Institute of Medical Sciences, Karad, Maharashtra, INDIA.
Email: ashars97@gmail.com

Received Date: 20/08/2021 Revised Date: 05/09/2021 Accepted Date: 12/10/2021

Access this article online

Quick Response Code:



Website:

www.statperson.com

Volume 11
Issue 4

INTRODUCTION

Ureterocele is dilatation of lower ureter at its entry into urinary bladder. It occurs due to incomplete dissolution of Chwallas' membrane. It is classified as single or duplex system and acquired or congenital. Acquired type is commonly seen in adults. It is associated with pathologies like impacted stone, schistosomiasis. Ureteral duplication anomalies are seen in 1 In 125 live births.¹ Duplicated ureters are often associated with an obstructed upper pole moiety and a refluxing lower pole moiety. Ectopic ureterocele typically arise from the upper pole moiety and are common in paediatric population.² Condition could be completely asymptomatic. The patient may present with pain radiating from flanks, associated with lower UTI problems, incontinence, urinary tract calculus and urethral prolapse. Diagnosis of ureterocele is confirmed

by its radiological appearance. Diagnosis may be made with ultrasonography in perinatal period or in early childhood. In adults intravenous urography demonstrates dilated distal ureter, appearing as a "cobra head" or "springonion" deformity with peripheral hollows. Renal functions are also confirmed on DMSA, DTPA or MAG³ scans. Adults presenting with symptoms of obstruction are treated to relieve obstruction, either endoscopically or open.³ Endoscopic incision and marsupialization would relieve the obstruction and preserve renal function.

CASE REPORT

18 year female presented with left flank pain radiating to groin. She did not have vomiting, hematuria or burning sensation during micturition. There was no similar complaint in the past. There was no previous surgery performed. There was no associated medical illness. No other person in the family presented with similar complaint. Her vitals were stable. Her abdominal examination revealed tenderness in right renal angle. External genitalia were normal on examination. Intravenous urogram performed revealed hydronephrosis and hydroureter of lower moiety of duplicated collecting system on left side. There was impacted calculus in ureterocele with another calculus in bladder. Kidneys were normally functioning on both sides. (Fig. 1) Her laboratory investigations were within normal range with normal serum urea and creatinine

values. The operation was performed to remove impacted calculus. Endoscopic incision and marsupialization was done for intravesicle ureterocele.



Figure 1: Radiograph showing ureterocele and hydronephrosis of lower moiety of duplicated draining system

DISCUSSION

Ureterocele is classified based on development as single system or duplex system. Mandell *et al* proposed a classification based on features of an affected ureteral orifice. American association of pediatricians has standardized terms, intravesical and ectopic according to extension into bladder, single or duplex system and in terms of orifices. Its incidence is 1:4000 children, common in Caucasians with female preponderance. Single system ureterocele is common in adults. It presents with other pathologies so it is also called as acquired type. In duplex system ureterocele, upper moiety is common.⁴ In our case report adult type simple ureterocele is presented in lower moiety of duplex system with impacted calculus. Intravenous urography showing complete duplication of left ureter both inserting in bladder. Lower moiety with hydronephrosis and hydronephrosis is seen. Ureterocele seen in the lower end of ureter with a calculus. Upper moiety is inserting into bladder normally. One more calculus is seen in the lumen of the urinary bladder. Since our case presented with calculi, it was treated with endoscopic incision and marsupialization of ureterocele with ureteroscopic retrieval of stone with checking patency of the ureters under fluoroscopic guidance. Decision regarding further treatment like re-implantation for reflux and nephrectomy in case of dysplastic or non-functioning kidneys is required. In case of re-implantation upper moiety is re-implanted in upper part and lower moiety is re-implanted in lower part of bladder. In our case renal function was intact and

obstruction was removed, no further intervention is required. Because of rarity of the presentation and diagnosis of the condition, urologist needs to be aware of etiology and system and in terms of orifices. Its incidence is 1:4000 children, common in Caucasians with female preponderance. Single system ureterocele is common in adults. It presents with other pathologies so it is also called as acquired type. In duplex system ureterocele, upper moiety is common.⁴ In our case report adult type simple ureterocele is presented in lower moiety of duplex system with impacted calculus. Intra venous urography showing complete duplication of left ureter both inserting in bladder. Lower moiety with hydronephrosis and hydronephrosis is seen. Ureterocele seen in the lower end of ureter with a calculus. Upper moiety is inserting into bladder normally. One more calculus is seen in the lumen of the urinary bladder.

Since our case presented with calculi, it was treated with endoscopic incision and marsupialization of ureterocele with ureteroscopic retrieval of stone with checking patency of the ureters under fluoroscopic guidance. Decision regarding further treatment like re-implantation for reflux and nephrectomy in case of dysplastic or non-functioning kidneys is required. In case of re-implantation upper moiety is re-implanted in upper part and lower moiety is re-implanted in lower part of bladder. In our case renal function was intact and obstruction was removed, no further intervention is required. Because of rarity of the presentation and diagnosis of the condition, urologist needs to be aware of etiology and appearance. Treatment options depend on the extension of the ureterocele, obstruction to draining system. Since few cases have been reported in literature, clinicians need to have understanding and orientation towards diagnosis, investigations and treatment of this condition.

REFERENCES

1. Schlussek, R.N., Retik, A.B. Ectopic ureter, ureterocele and other anomalies of the ureter. Campbell-Walsh Urology. 9th edn: p.3413
2. Donald S. Hill. 2004. Case studies, Prolapsed, ectopic ureterocele. Journal of Diagnostic Medical Sonography. 20(6):436-439
3. Byun E. Merguerian PA. 2006. A meta-analysis of surgical practice pattern in endoscopic management of ureteroceles. Journal of Urology. 176(4):1871-1877
4. Chtourou M, Sallami S, Rekik H, Binous *et al*. 2002. Ureterocele in adults complicated with calculi: Diagnostic and Therapeutic features. Reports of 20 cases. Prog Urol. 12:1213-1220.

Source of Support: None Declared
Conflict of Interest: None Declared