

Mucopolysaccharidosis type I-Hurler's syndrome

A M Deodhar¹, Vedprakash Biradar^{2*}, Rushikesh M Patwardhan³, Parikshit Mule⁴, Pallavi Rawal⁵, Rajeshri Ekale⁶

Vivekanand Hospital, Signal camp, Vidya nagar, Latur-413531, Maharashtra, INDIA.

Email:vedprakash1908@gmail.com

Abstract

Mucopolysaccharidosis I (MPS I) is a rare inherited disorder that belongs to a group of clinically progressive disorders and is caused by the deficiency of the lysosomal enzyme, $\alpha 1$ -iduronidase. MPS I has been recently classified into a severe (Hurler syndrome) and an attenuated type (Hurler-Scheie and Scheie syndromes).

Keyword: $\alpha 1$ -iduronidase, Hurler-Scheie syndrome.

*Address for Correspondence:

Dr. Vedprakash Biradar, Vivekanand Hospital, Signal camp, Vidya nagar, Latur-413531, Maharashtra, INDIA.

Email:vedprakash1908@gmail.com

Received Date: 22/08/2021 Revised Date: 04/09/2021 Accepted Date: 10/10/2021

Access this article online	
Quick Response Code:	Website: www.statperson.com
	Volume 11 Issue 4

lipoid accumulation in central nervous system and other viscera. Its incidence is 1:100000 of births. The excessive mucopolysaccharides excreted in urine are dermatan sulfate and heparan sulfate. Their synonym includes lipochondrodystrophy, Gargolism, osteochondrodystrophy, dysostosis multiplex.

CASE HISTORY

3years old male child presented with short stature (stunted growth), coarse facial features, limping and spinal deformity. Patient was subjected for radiograph of thoracic-lumbar spine AP and Lateral view, Both Hand AP view, skull AP and lateral view. Radiograph revealed

INTRODUCTION

Hurler's syndrome is rare autosomal recessive disorder of mucopolysaccharid metabolism that leads to excessive




Figure 1: Thoraco-lumbar AP and Lateral view-> kyphosis of spine, Lower thoracic (T9 to T12) and upper lumbar vertebrae shows anterior-inferior beaking, appears small in antero-superior aspect, rest visualized thoracic vertebrae appears oval in shape. Visualized lower ribs appear wide anteriorly producing spatulated appearances. Ilii are flared, with obliquely directed acetabular roof



Figure 2




Figure 3

Legend

Figure 2: Both Hands AP view-> Metacarpals and phalanges are short, wide producing trident hand. Bullet shaped metacarpals

Figure 3: Skull AP and lateral view-> Macrocephaly, frontal bossing, calvarial thickening

Findings suggestive of Mucopolysaccharidosis 1-Hurler's syndrome.

DISCUSSION

Mucopolysaccharidosis-I (MPS I) is a lysosomal storage disorder inherited as an autosomal-recessive condition and is caused by a deficiency of the lysosomal enzyme α 1-iduronidase. This results in the progressive accumulation of glycosaminoglycans (GAG) within the lysosomes, leading to multiorgan dysfunction and damage¹. Patients affected with MPS I are unable to degrade the GAG, dermatan sulfate, and heparan sulfate, which provide structural support to the extracellular matrix and cartilaginous structure such as joints and heart valves². Patients are usually normal at birth and remain so until after the first year of life. Facial features then begin to coarsen, with the development of large head, wide set eyes (hypertelorism), sunken nose, large lips and protruding tongue. Corneal opacities develop and teeth are short and malformed. Mental retardation, deafness gradually develops.

Hepatosplenomegaly, protruberant abdomen, umbilical and inguinal hernias are common. Eventually the patients become dwarfed. A severe dorso-lumbar kyphosis develops³. MPS I has been classified into two broader groups, severe MPS I (Hurler Syndrome) and attenuated MPS I (Hurler-Scheie and Scheie syndromes)⁵. The greatest variability is observed in individuals with the attenuated MPS I. Onset is usually between ages three and ten years. Although psychomotor development may be normal in early childhood, individuals with attenuated MPS I may have learning disabilities. The rate of disease progression and severity can range from serious life threatening complications (leading to death in the second to third decades) to a normal life span with significant disability and discomfort from progressive severe restriction in the

range of motion of all joints. Hearing loss and cardiac valvular disease are common. The diagnosis of MPS I relies on the demonstration of deficient activity of the lysosomal enzyme α -L-iduronidase in peripheral blood leukocytes, cultured fibroblasts, or plasma. Glycosaminoglycan (GAG) (heparan and dermatan sulphate) urinary excretion is a useful preliminary test⁶. Other radiological findings include -premature closure of sagittal and lambdoid sutures. Hydrocephalus is common. Sella turcica is enlarged and J shaped. Dens hypoplasia resulting into atlantoaxial subluxation. In Pelvis illi are flared with obliquely directed acetabular roof. Coxa vera or vulga is common. Varus deformity of humerus is characteristic⁷. On sonography hepatomegaly may be seen.

REFERENCES

1. Lamy M, Maroteaux P, Bader JP; Etude genetique, J Genet Hum 6:156,1957
2. Mikles M Stanton RP; A review of Hurlers syndrome Am J Orthop 25(8);533,1997
3. Thomas SL Tandon, Hurlers syndrome, A case report 24(2).245 2000
4. Kressler R J, Aergarter EE; Hurlers syndrome summary of literature and case report without autopsy findings. J Pediatr 12:576,1938.
5. Eggli KD, Dorst JP: Mucopolysaccharidosis and related condition 21;275,1986.
6. Neufeld EF, Muenzer J. The mucopolysaccharidoses. In Scriver C, Beaudet A, Sly W, et al, editors. *The metabolic and molecular bases of inherited disease*. McGraw Hill: New York, NY 2001; .3421-52.
7. Terry R. Yochum, Lindsay J. Rowes: Essentials of skeletal Radiology; 3rd edition, page 745.

Source of Support: None Declared
Conflict of Interest: None Declared