

Starch and Glucose Content and Changes therein in the Seeds of *Schleichera oleosa*(kusum) due to Bioteriation by Pathogenic Fungi during Storage.

AK Srivastava* and GK Pandey**

*Head, Dept of Botany, St .Xavier's College, Ranchi INDIA

**Dept of Botany, St Columba's PG College, Hazaribag, INDIA

Corresponding Address:

ajaysrivastava11@gmail.com

Research Article

Abstract:: Seeds of one of the non-edible oilseed, kusum(*Schleichera oleosa* f. *Sapindaceae*) are of great significance in Jharkhand area and its oil is used in burning lamps, varnishing, massage and medicine. Its oil-cake is a good manure. The unsanitary and humid condition makes it prone to a faster biodeterioration due to several fungal inhabitants during storage. The colonising fungi in the seeds use up some of the glucose and starch as carbon sources for meeting energy requirements in their prolonged association with the seeds. In the span of one year the glucose level dropped by as much as 50.2 percent of its initial value. The starch in the stored seeds reduced steadily from 2.55 percent of dry seed weight to 1.25 percent due to the association of the mycodwellers.

Keywords : *Schleichera oleosa*, oilseed, colonising fungi, carbon sources, mycodwellers.

Introduction:

Seed pathology is up and coming area of plant pathology and has attained a great magnitude. Seeds of a non edible oil, *Schleichera oleosa* f. *Sapindaceae* (Kusum) are of great use in burning lamps, varnishing, massage and medicine. Its oil-cake is good manure. In the Jharkhand area, Kusum seeds are collected just after onset of rainy season i.e. by the end of month of May. Conventionally seeds are stored in gunny bags in the village houses and godowns. In either case the condition is unsanitary and humid and is conducive for fungal growth leading to good scale destruction.

There are many reports indicating that fungi are associated with seed surface as well as internal mycoflora(DUTTA & ROY , 1987 ; JAYWEERA *et al* 1988). It was worthwhile to seek the extent of

biodeterioration occurring in the seeds in terms of carbohydrates in different forms viz. glucose and starches.

Material and Methods:

Kusum seeds were obtained from the godown of Ranchi forest department for the preparation of oil and oilcake on regular interval of one month. The work spanned from June 2001, one harvest season to May 2002, just before the other harvest season. Glucose content was estimated by titration method using Fehling solutions (Cole's method, 1933). Starches were estimated by colorimetric method using anthrone as a reagent (DUBOIS *et. al.*, 1951) using a colorimeter of Systronics (India) make. Estimations were done in triplicate and standard deviation and standard errors were calculated.

Results and Discussion:

The effect of continued fungal activity on the seeds manifested in the lowered glucose and starch contents just like oils (SRIVASTAVA, AK and PANDEY GK, 2000). The glucose level dropped by as much as 50.2 percent in the span of one year of its initial value. The starch in the stored seeds reduced steadily from 2.55 percent of dry seed weight to 1.25 percent due to the association of the mycodwellers. The results are elaborated in tables and figs 1-2.

Table 1: Amount of glucose present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack , June '01- May '02.

S No	Month	Wt.(mg)	Percentage	Change(mg)
1	Jun '01	16.75	1.675	-----
2	Jul '01	16.25	1.625	-0.05
3	Aug '01	16.25	1.625	-0.05
4	Sept '01	15.75	1.575	-0.1
5	Oct '01	14.9	1.49	-0.175
6	Nov '01	14.3	1.43	-0.245
7	Dec '01	13.1	1.31	-0.365
8	Jan '02	13.0	1.3	-0.375
9	Feb '02	11.9	1.19	-0.485
10	Mar '02	10.3	1.03	-0.645
11	Apr '02	9.5	0.95	-0.725
12	May '02	8.35	0.835	-0.84
	S.D. 0.284	S.E 0.081		

Final percent change in glucose content $[(1.675 - 0.835)/1.675] 100 = 50.2\%$




Fig 1: Amount of glucose present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack, June '01- May '02.

Table 2: Amount of starch present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack, June '2001 - May '02

S No	Month	Wt.(mg)	Percentage	Change(mg)
1	Jun '01	25.5	2.55	-----
2	Jul '01	24.0	2.4	-0.15
3	Aug '01	22.3	2.23	-0.22
4	Sept '01	22.0	2.2	-0.25
5	Oct '01	21.6	2.16	-0.39
6	Nov '01	19.9	1.99	-0.56
7	Dec '01	17.9	1.79	-0.76
8	Jan '02	17.1	1.71	-0.84
9	Feb '02	16.3	1.63	-0.92
10	Mar '02	14.8	1.48	-1.07
11	Apr '02	12.9	1.29	-1.26
12	May '02	12.5	1.25	-1.30
	SD:0.430	S.E.: 0.124		

Final percent change in starches content $[(25.5-12.5)/25.5]100 = 50.98\%$




Fig 2: Amount of starch present in 1 g of Kusum seed powder samples through one year of biodeterioration due to fungal attack , June '2001 - May '02

Changes in glucose content with pathogenic association have been reported. CHAUDHARY and PRASAD (1974) found a depletion of glucose among other nutrients in the shoot tissues of pigeon pea infected with *Fusarium oxysporum* f.sp.udum . CHATTOPADHYAYA and NANDY (1978) a carbohydrate loss to *F. subglutinum* infection. Loss in amount of glucose in fruits have been reported for tomato- *Drechslera australiense* (KAPOOR and TANDON, 1970); tomato-*Alternaria solani* (MEHTA et. al., 1975); banana-*Gloeosporium musarum*(WANG , 1960) and citrus-*Xanthomonas campestris* pv.citri (VIDYASEKARAN and DURAIRAJ, 1971) host pathogen systems.

Similarly the depletion in the starch content is commensurate with the activation of starch degrading enzymes especially **beta - amylase** (SCHIPPER and MIROCHA, 1977) . The drop is due to the sugar being used by the fungal pathogens as respiratory substrates(BAKER, 1965; WU , 1973) . In pigeon pea seeds infested with *Aspergillus flavus*, SINHA and PRASAD (1977) found a depletion of starches. Likewise BILGRAMI et.al., (1979 b) recorded a considerable decrease in the amount of starch in paddy seeds during 60 and 90 days of fungal infestation of an aflatoxin producing strain of *Aspergillus parasiticus*. SINHA et.al., (1981) found a considerable reduction in the starch content of *Cajanus cajan* seeds infested with *Aspergillus flavus* and *A. Niger*. However , these seeds , when infested with *Alternaria alternata* and *Curvularia lunata*, showed a moderate reduction in the starch contents . On the contrary, *Cajanus* seeds infested with *Fusarium moniliforme* and *Drechslera hawaiiensis* exhibited a minimum level reduction in starch contents. Later on, SINGH and SINHA (1985) confirmed that infestation of *Cajanus* seeds by *Aspergillus parasiticus* caused a considerable decline in their starch contents. PRASAD (1989) reported a loss of starch in fungi infested seeds of

Coriandrum indicum and the maximum loss in starch was due to *Aspergillus flavus* followed by *Curvularia lunata*. All these reports are concurrent to our findings. It is thus apperant that one year of seed infestation of *Schleichera oleosa* predominantly and jointly by *Aspergillus fumigatus*, *A. flavus*, *A. niger*, *Fusarium solanii*, *Paecilomyces variotii* and *Mucor sp.* causes biodeterioration of its seeds manifested in the diminishing of carbohydrate contents.

References:

- [1] Baker .J 1965. Study in the respiratory and carbohydrate metabolism of plant tissues.XVIII. The effect of oxygen on starch formation and dissolution in potatoes. New Phytol. 64: 201-209.
- [2] Bilgrami KS., Jamaluddin., Sinha RK. and Prasad T. 1979. Changes in seed contents of paddy(*Oryza sativa* L) due to fungal flora. *Phytopath J.* 96: 9- 14.
- [3] Chattopadhyay NC., and Nandy B.1978. Changes in nitrogen content in malformed inflorescence of mango caused by *Fusarium moniliforme* Sheld. Var subglutinans *Wr. Et. Rg Giornale Bot. Ital.* 112: 343-346.
- [4] Cole, SW.1919. The estimation of lactose and glucose by the copper *Biochem J.* 13(2).
- [5] Dubois ,M., Gilles,K., Hamilton , JK., Rebers,PA and Smith, F., 1951 A colorimetric method for the estimation of sugar. *Nature* 168: 167.
- [6] Kapoor , IJ and Tandon RN. 1970. Post infection changes in sugar content of tomato fruits caused by *Drechslera australiense*. *Indian Phytopath .* 23 : 133-135.
- [7] Mehta P., Vyas, KM and Saksena, SP. 1975. Metabolic changes during pathogenesis of fruit rot disease of tomato. *Indian Phytopath.* 28 : 253- 255.
- [8] Prasad M 1989 . Plant - bacteria interaction - Its dynamics and dimensions. In *Plant Microbe Interactions*. Ed: KS Bilgrami. Narendra Publishing House, Delhi : 143- 161.
- [9] Schipper , AL and Mirocha, CJ, 1977. Mechanism of starch hydrolysis in bean leaves during infection by bean rust fungus. *Phytopathology* 58 : 1066.

- [10] Singh , Premlata and Sinha KK. 1985. Changes in the seed content of arhar infected with *Aspergillus parasiticus*. *Indian Phytopathol.* **38** : 560.
- [11] Sinha MK and Prasad ,T . 1977. Deterioration of arhar seeds by *Aspergillus flavus*. *Indian Phytopathol.* **30** : 70-72.
- [12] Sinha ,MK., Singh, KK and Prasad , T. 1981. Changes in starch contents of Arhar seeds due to fungi. . *Indian Phytopathol.* **34** : 269- 271.
- [13] Srivastava ,AK and Pandey GK. 2000. Chemical changes in properties of kusum(*Schleichera oleosa*) oil during its seed infestation by fungi. *J.Mycopathol . Res.* **38(1)**: 29-32, (2000)
- [14] Wang, MC 1960. Physiological studies on *Gloeosporum musarum* Cook. At Mass, the causal organism of banana anthracnose . Changes in the carbohydrate composition of banana pulp with reference to the adaptive secretion of *amylase*. *Bot. Bull. Acad. Sin. N.S.* **1** : 59- 75.
- [15] WU , LC. 1973. Changes in some enzymes of moong bean seeds germinated on mycelia macerate of *Rhizoctonia solani*. *Physiol. Plant Pathol .* **3**: 19-27.