

Ultrasonographic correlation of abnormal uterine bleeding at perimenopausal stage

Jaideep M Palwade^{1*}, Charushila S Borole²

¹Assistant Professor, Department of OBGY, MIMER Medical College, Talegaon Dabhade, Pune, Maharashtra, INDIA.

Email: palwadejaideep@gmail.com

Abstract

Aims and Objectives: To study Ultrasonographic correlation of abnormal uterine bleeding at perimenopausal stage. **Methodology:** This was cross-sectional study carried out in 150 patients gynaecology OPD with the complaints of excessive per vaginal bleeding and who were in the age group between 37 to 51 years were considered. This study was carried out during year 2016. Total 150 females admitted in obstetrics and gynaecology department in MIMER Medical College were included into study. Clinical diagnosis and ultrasonographic findings were correlated. **Results:** Most of the patients in the present study were between 37-41 years i.e. 56%. Followed by 42-46 i.e. 25.33% and in 47-51 were 18.67 %. Majority of the patients on Ultrasonographic findings Showed no obvious pathology in 62 % patients, Leiomyoma in 21.33 %, followed by Adenomyosis in 5.33%, s/o PID in 2%, s/o Carcinoma growth in 2%. Respective percentage of Clinically diagnosed patients and Ultrasonographically diagnosed patients for Leiomyoma was 21.33 % , 21.33%, followed by DUB, PID, Carcinoma cervix was 2% , 2%; 7%, 3%; 44%, 7.33%. **Conclusion:** Except DUB, all other cases of AUB correlated well clinically and ultrasonographically. Leiomyoma and cervical growth were the only entities, which correlated well clinically, ultrasonographically. Whereas DUB, adenomyosis, polyp, PID did not correlate well clinically, ultrasonographically.. Adenomyosis was a histopathological diagnosis. DUB was an overestimated diagnosis clinically.

Key words: Abnormal uterine bleeding (AUB), Dysfunctional Uterine Bleeding(DUB) Perimenopausal stage.

*Address for Correspondence:

Dr. Jaideep M Palwade, Assistant Professor, Department of OBGY, MIMER Medical College, Talegaon Dabhade, Pune, Maharashtra, INDIA.

Email: palwadejaideep@gmail.com

Received Date: 06/11/2016 Revised Date: 16/12/2016 Accepted Date: 02/02/2017

Access this article online

Quick Response Code:



Website:
www.statperson.com

Volume 7
Issue 1

reproductive hormones can lead to an array of physical and psychological symptoms such as Changes in menstrual patterns, Vasomotor symptoms, Psychological and mental disturbances, Sexual dysfunction, Somatic symptoms , Dry, itchy skin.²

Abnormal uterine bleeding (AUB) is a symptom and not a disease. It is one of the most frequently encountered complaints in gynecologic practice. It accounts for more than 70% of all gynaecological consultations in the perimenopausal age group.³ It occurs in various forms such as menorrhagia, polymenorrhea, polymenorrhagia, metrorrhagia, and menometrorrhagia.⁴ The International Federation of Gynaecology and Obstetrics in November 2010, accepted a new classification system i.e. PALM-COEIN for causes of AUB in the reproductive years. The system based on the acronym (polyps, adenomyosis, leiomyoma, malignancy and hyperplasia-coagulopathy, ovulatory disorders, endometrial causes, iatrogenic, not classified) was developed in response to concerns about the design and interpretation of basic science and clinical investigation that relates to the problem of AUB.⁵ AUB may be an expression of hormonal milieu, or it could be the clinical presentation of benign or malignant lesions of female genital tract in perimenopausal woman. However,

INTRODUCTION

Different opinions are expressed as which years in women's life should be accepted as perimenopausal period. Perimenopause refer to the time period in the late reproductive years, usually late 40s to early 50s. Characteristically, it begins with menstrual cycle irregularity and extends to 1 year after permanent cessation of menses. The more correct terminology for this time is menopausal transition. This transition typically develops over a span of 4 to 7 years, and the average age at its onset is 47 years.¹ During the menopausal transition, more erratic fluctuations in female

there are no detectable structural abnormalities in majority of cases and this is called dysfunctional uterine bleeding (DUB). DUB, fibroid uterus, and adenomyosis are the common hyperoestrogenic conditions where endometrium remains in the proliferative phase and if untreated may lead to endometrial carcinoma. Therefore, clinical examination and investigations are essential to find out the etiological factor in a perimenopausal patient presenting with AUB. Ultrasonography (USG) can be used to exclude organic pathology for AUB. AUB is one of the main gynaecological reasons of hysterectomy and accounts for two-thirds of all hysterectomies.⁶

METHODOLOGY

This was cross-sectional study carried out in gynaecological OPD with the complaints of excessive per vaginal bleeding and who were in the age group between 37 to 51 years were considered. Also women complaining of abnormal uterine bleeding per vaginum in some form or other were studied. A detailed history of the patient was obtained taking into account any associated symptoms like dysmenorrhea, dyspareunia, postcoital bleeding, intermittent spotting, unhealthy discharge, foul smelling discharge, heaviness and discomfort in the lower abdomen, backache and any other constitutional symptoms was obtained. The study was carried out at over a period of 12 months from Jan 2016 to Dec 2016 in obstetrics and gynaecology department of MIMER Medical College, Talegaon Dabhade. In this period total 150 patients were included into study. Per Speculum Examination done to note the condition of vagina - any erosion, growth, suspicious looking area, any polyp seen, any discharge, Per Vaginal Examination - To note, Size and position of the uterus, Mobility, Consistency, Irregularity, Tenderness, Any adnexal lump, Per Rectal Examination- in all suspected cases of carcinoma, Routine investigations, Hemoglobin Urine routine / microscopy, Peripheral blood smear, Platelet count, Bleeding time, Clotting time, HIV test, HbsAg, Blood sugar random etc. done for confirmation of clinical

diagnosis and same patients underwent Ultrasonographic investigations. The findings were correlated.

RESULT

Table 1: Age Distribution of Women with Abnormal Uterine Bleeding At Perimenopausal Age

Sr. No.	Age Group of Patients in Years	No. of Patients	Percentage
1	37-41	84	56
2	42-46	38	25.33
3	47-51	28	18.67
Total		150	

Most of the patients in the present study were between 37-41 years i.e. 56%. Followed by 42-46 i.e. 25.33% and in 47-51 were 18.67%.

Table 2: Ultrasonographic Diagnosis

Sr. No.	Ultrasonographic Findings(n=150)	No. of cases (Percentage of Total Cases)
1	DUB	11 (7.33)
2	Leiomyoma	32 (21.33)
3	Adenomyosis	8 (5.33)
5	s/o PID	3 (2)
6	s/o Carcinoma growth	3 (2)
7	No obvious pathology	93 (62)
Total		150

(P<0.005)

From above Table it is clear that the majority of the patients on Ultrasonographic findings Showed No obvious pathology In 62 % patients, Leiomyoma in 21.33 % , followed by Adenomyosis in 5.33% , s/o PID in 2% , s/o Carcinoma growth in 2%.




Figure 1:

Table 3: Correlation between Clinical, Ultrasonographic, and Histopathological Examination

Sr. No.	No. of cases	Clinical Diagnosis No of Cases (Percentage)	USG Diagnosis No of Cases (Percentage)	HP Diagnosis after Sampling No of Cases (Percentage)
1	DUB	66 (44)	11 (7.33)	52 (34.67)
2	Leiomyoma	32 (21.33)	32 (21.33)	32 (21.33)
3	Adenomyosis	21 (14)	8 (5.33)	33 (22)
4	Polyp	18 (12)	0 (0)	18 (12)
5	PID	7 (4.67)	3 (2)	8 (5.33)
6	Carcinoma Cervix	3 (2)	3 (2)	3 (2)
7	Carcinoma Endometrium	0 (0)	0 (0)	1 (0.67)
8	Others	3(2)	0 (0)	-

Other category includes AUB due to thrombocytopenia and aspirin (anti-platelet drug).




Figure 2:

From above Table and Graph it is clear that maximum correlation in Clinical Ultrasonographic, Histopathological was found for Leiomyoma i.e. 21.33 % , 21.33%, 21.33% followed by Carcinoma of Cervix, PID, DUB i.e. 44%, 7.33%, 34.67%; 4.67, 2, 5.33; 44, 7.33, 34.67 etc.

DISCUSSION

The TVUS is a simple and non-invasive diagnostic modality of studying the endometrial pattern and its thickness accurately, and at the same time to exclude organic pathology in cases of AUB. The high frequency transducer placed nearer to the region of interest permits better visualisation of the uterus and the endometrium.¹⁰ In our study Findings suggestive of no obvious pathology were detected on ultrasonography in 62% of total 150 cases as shown in Table 10. Leiomyoma was the sonographic diagnosis in all clinically suspected myoma cases (100%). Ultrasonographic evidence of carcinoma cervix was observed in all 3 cases of carcinoma cervix. (100%). Ultrasonography was suggestive of adenomyosis in 38.1% of total clinically suspected adenomyosis cases. DUB was diagnosis in 11 cases. No obvious pathology is highly significant than the other ultrasonographic findings, calculated statistically which was <0.001 (Table 2). No obvious pathology included clinically diagnosed cases of PID, polyp, DUB, adenomyosis. Clinical as well as ultrasonography evaluation proved less useful for diagnosing adenomyosis and DUB. Leiomyoma was the sonographic diagnosis in all clinically suspected myoma cases (100%) in a study by Archana B *et al* (2010).¹¹ The final pathologic (true) diagnosis confirmed the clinical indication in all cases (100% correlation) of leiomyomas, adenomyosis and endometrial polyps.¹² A study was conducted on terminology and evaluation of abnormal uterine bleeding in perimenopausal women, evaluated a sample of 433 perimenopausal patients with abnormal uterine bleeding. Out of 341 patients, 79% had ultrasonographic evidence of no anatomic abnormality.¹³ These findings are in confirmation with study, Bharat Talukdar¹⁴ 103 numbers of perimenopausal hysterectomized patients were analyzed. Most number of patients (69.67%) were between 40 and 45 years age group. The common menstrual problem was menorrhagia

(43.69%). This finding was comparable with the study of Jetley *et al*.⁷ and Shobha,⁸ in which clinical presentation as menorrhagia in AUB evaluation revealed 46.4% and 46.6%, respectively. It appeared in this study that the maximum patients attended hospital for treatment after suffering for 3–6 months (54.37%) which was comparable to the study of Kathuria and Bhatnagar (50%).⁹ also in confirmation with Shobha S. Pilla¹⁵.

CONCLUSION

Except DUB, all other cases of AUB correlated well clinically and ultrasonographically. Leiomyoma and cervical growth were the only entities, which correlated well clinically, ultrasonographically. Leiomyoma, cervical growth correlated well clinically and ultrasonographically. Whereas DUB, adenomyosis, polyp, PID did not correlate well clinically, ultrasonographically. Adenomyosis was a histopathological diagnosis. DUB was an overestimated diagnosis clinically.

REFERENCES

1. McKinlay SM, Brambilla DJ, Posner JG. William's gynecology. New York: McGraw-Hill. 2008; 468-491.
2. Bachmann G. Physiologic aspects of natural and surgical menopause. J Reprod Med. 2001; 46(3):307.
3. Mahajan N, Aggarwal M, Bagga A. Health issues of menopausal women in North India. J Midlife Health. 2012; 3:84-7.
4. Kumar P, Malhotra N. Clinical types of abnormal uterine bleeding. In: Kumar P, editor. Jeffcoate's Principle of Gynecology. 7th ed. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd; 2008. p. 599.
5. Munro MG, Critchley HO, Fraser IS. The FIGO systems for nomenclature and classification of causes of abnormal uterine bleeding in the reproductive years: Who needs them? Am J ObstetGynecol. 2012; 207:259-65.

6. Telner DE, Jakubovicz D. Approach to diagnosis and management of abnormal uterine bleeding. *Can Fam Physician*. 2007; 53:58–64.
7. Jetley S, Rana S, Jairajpuri ZS. Morphological spectrum of endometrial pathology in middle-aged women with atypical uterine bleeding: A study of 219 cases. *J Midlife Health*. 2013; 4:216–20.
8. Shobha PS. Sonographic and histopathological correlation and evaluation of endometrium in perimenopausal women with abnormal uterine bleeding. *Int J Reprod Contracept ObstetGynaecol*. 2014; 3:113–7.
9. Kathuria R, Bhatnagar B. Correlation between DandC, USG and hysteroscopy findings in diagnosing a cause for abnormal uterine bleeding. *Indian J ClinPract*. 2014; 25:466–70.
10. Goldstein SR. The endometrial echo revisited: have we created a monster? *Am J ObstetGynecol*. 2004; 191:1092-6.
11. Archana B, Michelle F. Evaluation and Histopathological Correlation of Abnormal Uterine Bleeding in Perimenopausal Women. *Bombay Hospital Journal*. 2010; 52(1):69-72.
12. Gupta G, Kotasthane DS, Kotasthane VD. Hysterectomy: A Clinico-Pathological Correlation of 500 Cases. *The InternandJournal of Gynecology and Obstetrics*. 2010; 14(1):145-152.
13. Pilli GS andal .Dysfunctional uterine bleeding. *The Journal of Obstetrics and Gynecology of India*. 2002 May-Jun; 52(3):87-9.
14. Bharat Talukdar , SangitaMahela J Midlife Health. Abnormal uterine bleeding in perimenopausal women: Correlation with sonographic findings and histopathological examination of hysterectomy specimens. *2016 Apr-Jun; 7(2): 73–77.*
15. Shobha S. Pillai. Sonographic and histopathological correlation and evaluation of endometrium in perimenopausal women with abnormal uterine bleeding. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology Pillai SS. Int J Reprod Contracept ObstandGynecol*. 2014 Mar;3(1):113-117.

Source of Support: None Declared

Conflict of Interest: None Declared