

The study of incidence of maternal anaemia in various age groups of women going to labour in Orissa

Rajeshwari K.^{1*}, Ashok Kumar Behera²

¹Senior Resident, Department of Obstetrics and Gynaecology, Chamarajanagara Institute of Medical Sciences, Chamarajanagara District, Karnataka, INDIA.

²Professor and HOD, Department of Obstetrics and Gynaecology, S C B Medical College, Cuttack, Orissa, INDIA.

Email: rajik09@yahoo.co.in

Abstract

Introduction: Anaemia is the commonest medical disorder in pregnancy and has a varied prevalence, aetiology and degree of severity in different populations. Anaemia during pregnancy has been shown to be associated with a two-fold risk for preterm delivery and a three-fold risk for low birth-weight as well as maternal mortality. The World Health Organization (WHO) estimates that anaemia contributed to approximately 20% of the 515,000 maternal deaths worldwide in 1995. Keeping these facts in view, the present study embodies the observation of 250 cases among 400 cases attending labour room of S. C. B. Medical College, Cuttack (Orissa), giving an incidence of 62.5%, which is quite high in comparison to developed countries.

Keywords: Maternal anaemia, incidence, age group.

*Address for Correspondence:

Dr Rajeshwari K., Senior Resident, Department of Obstetrics and Gynaecology, Chamarajanagara Institute of Medical Sciences, Chamarajanagara district, Karnataka, INDIA.

Email: rajik09@yahoo.co.in

Received Date: 17/04/2018 Accepted Date: 27/06/2018

Access this article online	
Quick Response Code:	Website: www.statperson.com
	Volume 8 Issue 3

INTRODUCTION

Anaemia is the commonest medical disorder in pregnancy and has a varied prevalence, aetiology and degree of severity in different populations¹. Anaemia in pregnancy is defined as a condition of low circulating haemoglobin in which the haemoglobin concentration has fallen below a threshold lying at two standard deviations below the median of a healthy population of the same age, sex and stage of pregnancy². WHO definition for diagnosis of anaemia in pregnancy is a haemoglobin concentration of less than 11 g/dl (7.5mmol/l) and a haematocrit of less than 0.33³. The prevalence of iron-deficiency anaemia in pregnant women is estimated to be between 35 and 75% (average 56%) in developing countries whereas in

industrialized countries the average prevalence is 18%^{4,5}. Out of an estimated 150 million deliveries occurring annually in the world, approximately 600,000 women die from complications of pregnancy and child birth^{6,7}. Anaemia is responsible for 40 – 60% of maternal death in non- industrialised countries. It causes direct as well as indirect, deaths from cardiac failure, hemorrhage, infection and pre-eclampsia^{8,9}. It also increases perinatal mortality and morbidity rates consequent to preterm deliveries, intra-uterine growth retardation, low iron stores, iron deficiency anaemia and cognitive and affective dysfunction in the infant^{10,11}. A number of studies have been done previously on the incidence of maternal anaemia in various age groups Malhotra *et al* (2002)¹², Aimakhu *et al* (2003)¹³. Keeping these facts in view the present study was conducted in this tertiary care hospital as cases from all the strata of the society come here. The study had the objective of finding out the magnitude of the problem in this part of the country.

MATERIAL AND METHOD

Source of Data

The present study was carried out in the department of Obstetrics and Gynaecology, SCB Medical College Hospital, Cuttack from 2009 to 2011.

Inclusion Criteria

Patients in labour with haemoglobin level of less than 11.0 gm / dl.

Exclusion Criteria

- Patients with haemoglobinopathies.
- Patients with ante-partum haemorrhage, bleeding disorder
- Pregnancy with bone marrow insufficiency
- Pregnancy with severe infections
- Grand multipara

Method of Study

A cross sectional study was conducted on women in labour with Hb <11gm/dl. All patients admitted in labour room had undergone haemoglobin estimation and women with Hb <11gm/dl were recruited in the study after they satisfied the inclusion and exclusion criteria. The age of women was recorded. The written informed consent was taken.

OBSERVATION AND RESULT

In the present study conducted on a sample of 400 cases attending labour room, percentage of incidence of anaemia on the whole and in various age groups are shown below in tables and bar diagram.

Table 1: Incidence of maternal anaemia

No. of Cases	No. of maternal anaemia Cases	Percentage
400	250	62.5

Above table shows detection of 250 maternal anaemia cases from a sample of 400 cases attending labour room, giving an incidence of 62.5%.

Table 2: Incidence of maternal anaemia in various age groups

Age in years	No. of Cases	Percentage
16 – 20	14	6
21 – 25	127	51
26 – 30	94	37
> 30	15	6
Total	250	100




Figure 1: Incidence of maternal anaemia among various age groups

Out of 250 cases, 14 cases (6 %) were between 16 – 20 yrs of age, 127 cases (51%) were between 21 – 25 yrs of age, 94 cases (37%) were between 26 – 30 yrs of age and 15 cases (6%) were above 30 yrs.

DISCUSSION

The present study was proposed to find out the incidence of anaemia in this part of the country in women going to labour. The present study embodies the observation of 250 cases of maternal anaemia among 400 cases attending labour room of S. C. B. Medical College, giving an incidence of 62.5% (Table I), which is quite high in comparison to developed countries. The study includes only cases that were anaemic at the onset of labour as it would have been unethical not to treat the cases to observe the effect of anaemia on pregnancy outcome. Adinma *et al* (2002)¹⁴ reported the incidence to be 17.2% at the onset of labour. Aimakhu *et al* (2003)¹³ reported an incidence of 15%. Sharma *et al* (2003)¹⁵ reported the prevalence of anaemia to be 96%. Dairo *et al* (2004)¹⁶ reported the prevalence of anaemia to be 32.8%. Rusia *et al* (2005)¹⁷ reported an incidence of 43.2%. Owusu *et al* (2005)¹⁸ reported the prevalence of moderate anaemia to be 57.15%. Singh *et al* (2006)¹⁹ reported an incidence of 84.9%. Anorlu *et al* (2006)²⁰ showed the prevalence as 35.2%. Marahatta (2007)²¹ reported the prevalence of anaemia to be 42.6%. Jaleel and Khan (2008)²² reported an incidence of 69.9%. Bukar *et al* (2008)²³ reported the prevalence to be 51.8%. Rasheed *et al* (2008)²⁴ reported an incidence of 41.3%. Noronha *et al* (2010)²⁵ reported the prevalence of 53.7%. Zugu Mei (2011)²⁶ reported prevalence to be 18±1.4% The prevalence of anaemia in pregnant women in different parts of India is given below^{27,28,29,30}

Table 3

Place	No. of pregnant women	Prevalence of anaemia
Assam	525	91.4%
Bihar	446	84.1%
U. P	593	79.4%
H. P	507	61%
J and K	498	96.8%
Tamil Nadu	1032	68.8%
Punjab	4752	86.1%

In the present study the incidence is comparable to the study of Jaleel and Khan (2008)²². The higher incidence of anaemia observed by the present study might be because most cases are referred from rural areas where anaemia antedates pregnancy and is aggravated by increased requirements during pregnancy. While studying the incidence of anaemia in various age groups (Table II) out of 250 cases maximum number of cases (87 %) belonged to the age group of 21 – 30 yrs. The increased incidence in this age group could be because of the fact that maximum number of patients do deliver in this age group. The youngest patient was of 19 years old and the oldest was 35 years old. The mean age was 25.5 years. Malhotra *et al* (2002)¹² reported the mean age as 27 ± 4.25. Aimakhu *et al* (2003)¹³ reported that 83.4% were

aged between 21 – 35 years. Sharma *et al* (2003)¹⁵ reported that the average age of the patient was 26.5 years, which is comparable to the present study.

CONCLUSION

Nutritional deficiency anaemia during pregnancy continues to be a major health problem in all non – industrialised countries, contributing significantly to high maternal and perinatal mortality and morbidity rates. India continues to be one of the countries with very high prevalence. National Family Health Survey (NFHS-3) reveals the prevalence of anaemia to be 70-80% in children, 70% in pregnant women and 24% in adult men. The present study showed an incidence of 62.5% with maximum number of cases (87 %) belonging to the age group of 21 – 30 yrs, which is quite high in comparison to developed countries. The high risk patients should be identified early and should be advised to have regular ANC and prophylactic iron and folic acid supplementation. Anaemia is a preventable condition, so all pregnant women must be observed and managed with adequate maternal and neonatal intensive care facilities to improve the outcome.

REFERENCES

- Schwartz WJ, Thurnau GR. Iron Deficiency Anaemia In Pregnancy. Clin Obstet Gynecol 1995; 38: 443- 454
- WHO/UNICEF/UNU. Indicators for assessing iron deficiency and strategies for its prevention: WHO draft, Geneva. WHO , 1996
- WHO, Report of WHO group of experts on Nutritional Anaemias. Technical report series no 503. Geneva WHO1972.
- World Health Organization. United Nations Children's Fund UNU. Iron deficiency anemia; Assessment, Prevention and Control; A guide for programme managers. World Health Organization, Geneva, 2001.
- Geneva WHO1DWNM922. The Prevalence of Anaemia in Women. WHO 1992; 1-99.
- WHO. Revised 1990. Estimates of Maternal Mortality. WHO/FRH/MSM/96.1. Geneva: WHO, 1996
- Turmen T, Abou Zahr C. Safe Motherhood. Int J Gynecol Obstet 1994; 46: 145 – 153.
- Bhatt R. Maternal mortality in India, FOGSI-WHO study. J. Obstet Gynecol India 1997; 47: 207-214.
- Viteri FE. The consequences of iron deficiency and anaemia in pregnancy. Adv Exp Med Biol 1994; 352 : 127 – 139
- Prema K, Neela KS, Ramalakshmi BA. Anaemia and adverse obstetric outcome. Nutri Rep Int 1981; 23: 637 – 643
- Lozoff B, Jimenez E, Wolf AW. Long term developmental outcome of infants with iron deficiency. N Engl J Med 1992 ; 325: 687 – 694
- Monika Malhotra, J B Sharma, S Batra, S. Sharma, N.S. Murthy, R. Arora, International Journal Of Gynecology and Obstetrics, Volume 79, Issue 2, Nov. 2002, pages 93-100
- CO Aimakhu, O Olayemi. West African Journal of Medicine > Vol 22, No 1 (2003)
- J I Brian-D Adinma, Joseph I Ikechebelu, UN Onyejimbe, G Amilo, Echendu Adinma. Tropical journal of Obstetrics and Gynecology, Vol 19, No 2 (2002)
- Jai Bhagwan Sharma: Dimple Soni: Nandagudi Srinivasa Murthy: Monika Malhotra. Department of Obstetrics and Gynecology, Department of Biostatistics, Maulana Azad Medical College, New Delhi, India .Journal of Obstetrics and Gynaecology Research, Volume 29, Issue 2, pages 73–78, April 2003
- Dairo MD, Lawoyin TO, Afr J Med Med Sci. 2004 Sep; 33(3):213-7.
- U. Rusia, C. Flowers, N. Madan, N. Agarwal, S. K. Sood and M. Sikka. Annals of Hematology . 2005, Volume 78, Number 8, 358-363, DOI: 10.1007/s002770050529
- Mary Glover-Amengor, W B Owusu, and BD Akanmori, Ghana Med J. 2005 September; 39(3):102–107.PMCID:PMC1790823.
- Toteja GS, Singh P, Dhillon BS, Saxena BN, Ahmed FU, Singh RP, Prakash B, Vijayaraghavan K, Singh Y, Rauf A, Sarma UC, Gandhi S, Behl L, Mukherjee K, Swami SS, Meru V, Chandra P, Chandrawati, Mohan U. Food Nutr Bull. 2006 Dec; 27(4):311-5.
- Anorlu RI, Oluwole AA, Abudu OO. J Obstet Gynaecol. 2006 Nov; 26(8):773-6.
- Marahatta, R, Nepal Medical College Journal. 2007 Dec; 9(4): 270-4.
- Riffat Jaleel, Ayesha Khan, Journal Of Surgery Pakistan (International) 13(4) Oct- Dec 2008
- M. Bukar , B. M. Audu, U. R. Yahaya and G. S. Melah. Anaemia in pregnancy at booking in Gombe, North-eastern Nigeria 2008, Vol. 28, No. 8 , Pages 775-778 (doi:10.1080/01443610802463835)
- Parveen Rasheed, Manal R Koura, Badria K Al-Dabal, Suhair M Makki Department of Family and Community Medicine, College of Medicine, King Faisal University, Dammam, Saudi Arabia, date of web publication 24-Aug-2009 Ann Saudi Med [serial online] 2008 [cited 2011 Jun 28];28:449-52.
- Maternal risk factors and anaemia in pregnancy: A prospective retrospective cohort study. J. A. Noronha, A. Bhaduri, H. Vinod Bhat, A. Kamath. February 2010, Vol. 30, No. 2, Pages 132-136.
- Zuguo Mei, Mary E Cogswell, Anne C Looker, Christine M Pfeiffer, Sarah E Cusick, David A Lacher, and Laurence M Grummer-Strawn. 2011 American Society for Nutrition
- Abel R, Rajarathnam J, Sampathkumar V. Anaemia in pregnancy. Impact of iron, deworming and IEC, RUSHA Dept. Tamilnadu: CMC Vellore 1999
- Indian Council of Medical Research. Evaluation of the National Nutritional Anaemia Prophylaxis Programme. Task Force Study. New Delhi: ICMR, 1989
- Sarin AR. Severe anaemia of pregnancy, recent experience. Int J Gynecol Obstet 1995; 50 (Suppl. 2): S45 – S49.
- Brabin L, Nicholas S, Gogate A, Karande A. High prevalence of anaemia among women in Mumbai, India. Food Nutr Bull 1998; 19: 205 – 209.

Source of Support: None Declared
Conflict of Interest: None Declared