

# Laparoscopy assisted management of vaginal evisceration

Jayanth Leo<sup>1\*</sup>, Parimuthukumar<sup>2</sup>, Vishwanath Pai<sup>3</sup>, A. Rekha<sup>4</sup>, Bhuvana<sup>5</sup>

{<sup>1,2</sup>Assistant Professor, <sup>3,4</sup>Professor, Department of General Surgery} {<sup>5</sup>Associate Professor, Department of Obstetrics and Gynaecology}  
Porur, Chennai 600116, Tamil Nadu, INDIA.

Email: [jayanthxl@yahoo.com](mailto:jayanthxl@yahoo.com)

## Abstract

**Introduction:** Evisceration is an uncommon surgical emergency. A vaginal evisceration is a rare occurrence following hysterectomy. The risk factors of having an evisceration are obesity, sudden increase in intra abdominal pressure, vaginal surgeries and post menopausal age group. This is a case report of a 58 year old lady who presented to the emergency room with evisceration through the vaginal vault. This mandates an early diagnosis of the condition, start the patient on broad spectrum antibiotics on admission and plan emergent surgical intervention. In our case we did a combined vaginal and laparoscopic approach. The use of minimally invasive surgery had saved the patient the morbidity of a laparotomy especially when done as an emergency procedure. The outcome of having managed this patient this way was rewarding.

**Keywords:** Evisceration, risk factors, hysterectomy, laparoscopy.

## \*Address for Correspondence:

Dr. Jayanth Leo, Department of General Surgery, Porur, Chennai 600116, Tamil Nadu, INDIA.

Email: [jayanthxl@yahoo.com](mailto:jayanthxl@yahoo.com)

Received Date: 08/02/2019 Accepted Date: 12/04/2019

Access this article online	
Quick Response Code:	Website: <a href="http://www.statperson.com">www.statperson.com</a>
	Volume 9 Issue 2

## INTRODUCTION

Vaginal evisceration was described by Hyernaux as early as 1864 as a disruption of the wall of proximal vagina resulting in a prolapse of abdominal contents. Most common organ to eviscerate is the small bowel. The etiopathogenesis is different for pre and post menopausal women with majority of cases having multiple risk factors aiding evisceration. There is no adequate data to understand the risk of vaginal vault dehiscence and there is no consensus on the management of this condition. It is important to identify and address these risk factors for a better understanding and complete management of the condition. The management of evisceration has to be tailored to individual patient to achieve the best outcome.

## CASE REPORT

A 58yr old woman was brought to the emergency room with prolapsing bowel loops through the vaginal orifice following alleged history of electric shock [fig 1]. She is a diabetic on medication and gives a past surgical history of abdominal hysterectomy done 6 years ago. On examination she was conscious, oriented and haemodynamically stable. Her abdomen was soft; however there was tenderness in the subumbilical region. Vaginal examination showed complete prolapse of vaginal vault with vault rupture and evisceration of small bowel loops. The bowel loops were grossly edematous. She was started on broad spectrum antibiotics with anaerobic cover. The eviscerated loops of intestines were promptly wrapped with warm saline soaks. In the operating room patient was under general anesthesia, in head low and lithotomy position. She was planned for Laparoscopy guided reposition of bowel loop along with vault repair. The eviscerated bowel loops, which was about 60 cms in length, were thoroughly washed with warm saline. There was a 2x1cm perforation in one of the bowel loop, which was closed in two layers using polyglactine sutures. An assistant all through the procedure to prevent further edema always supported the bowel loops. Laparoscopic ports were introduced, only to find dense bowel adhesions to the anterior abdominal wall (previous scar) in the lower segment of the abdomen.

This made visualization of the pelvis difficult. Anticipating such a problem the first port was made higher up than usual in the supra umbilical region and with a steep head low position [fig 2]. Acceptable exposure of the pelvis was attained and we managed to reduce the eviscerated bowel intraperitoneally by using the combined vaginal and laparoscopic approach. The entire length of the eviscerated bowel was placed back

intraperitoneally after which the two rents in the vaginal vault [fig 3] were closed with absorbable continuous interlocking sutures [fig 4]. The vault was repositioned with a vaginal pack insitu. Patient was shifted to ICU for overnight observation and management. She was subsequently shifted to ward and discharged on the fourth POD.



**Figure 1:** Clinical presentation. Complete vault prolapse with eviscerated edematous small bowel loops



**Figure 2:** Introduction of laparoscope



**Figure 3:** Rent in the vault seen after repositioning the bowel loops intraperitoneally



**Figure 4:** Completed vault closure

## DISCUSSION

Vaginal evisceration is not a commonly occurring phenomenon. It is rare and potentially life threatening. The actual incidence of such an event is difficult to ascertain. It is because of the rarity of such a complication, the incidence is low and is difficult to study. In an expert review by Cronin, the rate of vaginal vault rupture ranges from 0.14 to 4.1%<sup>1</sup>. This is however dependent on the type of surgical approach and patient factors. The risk factors of vaginal evisceration are different among pre and post menopausal age groups. Post menopausal women are more vulnerable to have evisceration because of the fact that there may be a preexisting prolapse of the vagina or the vaginal vault following hysterectomy<sup>2,3</sup>. Data on risk factors are limited and conflicting<sup>1</sup>. Even though it may be scientifically interpreted that age and medical conditions like diabetes may interfere with wound healing, there is no data available to substantiate this view. Elderly women, vaginal surgeries, vaginal atrophy, factors with poor wound healing (including malignancy, chronic steroid

use, malnutrition, tissue radiation), increased intra abdominal pressure (chronic cough and constipation), and postoperative vaginal cuff infection or hematoma may be risk factors favoring dehiscence<sup>4</sup>. The timing of dehiscence after hysterectomy is also very variable, it is reported as early as 3 days<sup>5</sup> to as late as 30 years<sup>6</sup> following surgery. So the vault giving way 6 years after surgery in this case is not surprising. There is no data associating obesity as a risk factor for having an evisceration. It is logical to accept however that the intra abdominal pressure in obese individuals is much higher as compared to the normal weight category<sup>7</sup>. This constant increased intra abdominal pressure along with the sudden increase in pressure precipitated in this patient by a fall on her abdomen following the electric shock, was one of the major factors responsible for the evisceration. The associated mortality is around 5.6% which gets worse if the eviscerated bowel is gangrenous<sup>10</sup>. It is of primary importance to assess and diagnose the condition early and plan adequate measures to restore anatomical integrity. It's always expected that

the eviscerated bowel loops are edematous and appear dusky due to venous congestion. This period is very vital to save the bowel from undergoing ischemic changes due to the increasing venous congestion<sup>8</sup>. Early administration of prophylactic antibiotics to control septicemia. The bowel when pushed intra abdominally carries the risk of contaminating the peritoneal cavity, especially since the bowel has been lying in contact with vaginal mucosa and the exterior. The basis of any surgical intervention offered to the patient aims at restoring the bowel back intra abdominally and closing the vaginal rent. In a review of case reports, series and retrospective studies done by Cronin and colleagues observed that a majority of cases that were reported were managed transvaginally and abdominally<sup>1</sup>. The author had taken into account all the available data for the last 30 years and concluded his review. We believe that use of laparoscopic assistance has been low since minimally invasive surgery has significantly matured only in the past decade. In this case, considering the laparoscopic findings in our patient (i. e. extensive bowel adhesions) on retrospect, one might favor a laparotomy over laparoscopy for better exposure and surgical ease. However with good optics and precise positioning during surgery this was overcome<sup>9</sup>. Also keeping in mind the patients age, BMI (47), nutritional status (S. albumin - 2.4) laparotomy might have been more morbid as compared to laparoscopy. The usage of laparoscopy has possibly averted grave wound complications following laparotomy.

## CONCLUSION

Vaginal evisceration is a rare but life threatening condition. It requires prompt diagnosis and early surgical intervention with equal emphasis on the use of prophylactic broad spectrum antibiotics. Combined

vaginal and laparoscopic approach is an effective method to reposition the bowel intraperitoneally<sup>10</sup>.

## REFERENCES

1. Beth Cronin, MD; Vivian W. Sung, MD, MPH; Kristen A. Matteson, MD, MPH; Vaginal cuff dehiscence: risk factors and management; American Journal of Obstetrics and Gynecology APRIL 2012
2. Kowalski CD, Seski JC, Timmins PF, Kanbour AI, Kunschner AJ, Kanbour-Shakir A. Vaginal Evisceration: presentation and management in postmenopausal women. J Am Coll Surg. 1996; 183:225-9.
3. Joy SD, Phelan M, McNeill HW. Postcoital vaginal cuff rupture 10 months after a total vaginal hysterectomy. A case report. J Reprod med. 2002; 47: 238-40.
4. Ramirez PT, Klemer DP. Vaginal evisceration after hysterectomy: a literature review. Ob- stet Gynecol Surv 2002; 57:462-7.
5. Cardosi RJ, Hoffman MS, Roberts WS, Spellacy WN. Vaginal evisceration after hyster- ectomy in premenopausal women. Obstet Gy- necol 1999; 94:859.
6. Moen MD, Desai M, Sulkowski R. Vaginal evisceration managed by transvaginal bowel resection and vaginal repair. Int Urogynecol J Pelvic Floor Dysfunct 2003; 14:218-20.
7. Wilson A, Longhi J, Goldman C, McNatt S; Intra-abdominal pressure and the morbidly obese patients: the effect of body mass index; J Trauma. 2010 Jul; 69(1):78-83. doi: 10.1097/TA.0b013e3181e05a79
8. Anthony Gyang, Antonio J. Ramirez, Spontaneous Vaginal Evisceration with Small Bowel Infarction; International Journal of Gynecology, Obstetrics and Neonatal Care, 2014, 1, 6-8
9. Yaakovian MD, Hamad GG, Guido RS. Laparoscopic management of vaginal evisceration: case report and review of the literature. J Minim Invasive Gynecol. 2008 Jan-Feb; 15(1):119-21. doi: 10.1016/j.jmig.2007.08.618.
10. Ioannis Nikolopoulos, Hasan Khan, Gnananandan Janakan, Rajab Kerwat; Laparoscopically assisted repair of vaginal evisceration after hysterectomy; BMJ Case Reports 2013; doi:10.1136/bcr-2013-009897

Source of Support: None Declared  
Conflict of Interest: None Declared